

Spectral dimension and diffusion in multi-scale spacetimes

Gianluca Calcagni¹ and Giuseppe Nardelli^{2,3}

¹ Instituto de Estructura de la Materia, CSIC, Serrano 121, 28006 Madrid, Spain

² Dipartimento di Matematica e Fisica, Università Cattolica, via Musei 41, 25121 Brescia, Italy

³ INFN Gruppo Collegato di Trento, Università di Trento, 38100 Povo (Trento), Italy

E-mail: calcagni@iem.cfm.csic.es, nardelli@dmf.unicatt.it

Abstract. Starting from a classical-mechanics stochastic model encoded in a Langevin equation, we derive the natural diffusion equation associated with three classes of multi-scale spacetimes (with weighted, ordinary, and ‘ q ’ Poincaré symmetries). As a consistency check, the same result is obtained by inspecting the propagation of a quantum-mechanical particle in a disordered environment. The solution of the diffusion equation displays a time-dependent diffusion coefficient and represents a probabilistic process, classified according to the statistics of the noise in the Langevin equation. We thus illustrate, also with pictorial aids, how spacetime geometries can be more completely catalogued not only through their Hausdorff and spectral dimension, but also by a stochastic process. The spectral dimension of multi-fractional spacetimes is then computed and compared with what found in previous studies, where a diffusion equation with some open issues was assumed rather than derived. These issues are here discussed and solved, pointing in favour of the model with q -Poincaré symmetries.

PACS numbers: 02.50.-r, 04.60.-m, 05.10.Gg, 05.40.-a, 05.45.Df

Contents

1	Introduction	2
2	Brownian motion, diffusion equation and spectral dimension	4
2.1	Brownian motion	5
2.2	Diffusion equation	6
2.3	Spectral and walk dimension	8
3	Multi-scale spacetimes	10
3.1	Laplacians	11
3.1.1	Weighted Poincaré symmetries	12
3.1.2	Ordinary Poincaré symmetries	12
3.1.3	q -Poincaré symmetries	13

3.2	Previous <i>Ansatz</i> for the diffusion equation	13
4	Diffusion in fractional spacetimes	16
4.1	Weighted Laplacian	16
4.1.1	Factorizable-spacetime Brownian motions	18
4.1.2	Spectral and walk dimension	22
4.2	Ordinary Laplacian	23
4.3	q -Laplacian	25
5	Diffusion in multi-fractional spacetimes	27
5.1	Weighted Laplacian	28
5.2	Ordinary Laplacian	30
5.3	q -Laplacian	31
6	Discussion	31
Appendix A Quantum probability density as a bilinear		33
Appendix B Quantum-mechanical time		35
B.1	Weighted Lagrangian	36
B.2	Ordinary Lagrangian	36
B.3	q -Lagrangian	37

1. Introduction

The concept of *spectral* (or *fracton*) *dimension* d_S of spacetime descends from the analogous one for a set, be it a fractal or a Riemannian manifold [1, 2, 3, 4, 5, 6]. The idea is to have a test particle diffuse in the set (a non-relativistic ‘space’), and see how it behaves. This process is governed by a diffusion equation, which for a Brownian motion reads

$$(\partial_t - \kappa_1 \nabla_x^2)P(x, x', t) = 0, \quad P(x, x', 0) = \delta(x - x'), \quad (1)$$

where the first-order derivative ∂_t is the diffusion operator in time t , κ_1 is a constant, ∇_x^2 (called spatial generator in probability theory) is the natural Laplacian in the given space (eventually containing metric structure) acting on the x dependence of the solution P , and x' is the initial point where the probe, a pointwise particle represented by the δ initial condition at $t = 0$, starts diffusing. The process has a probabilistic interpretation when P is a probability density function (PDF), $P \geq 0$ for all x . In particular, the random position field $X(t)$ associated with equation (1) is a Brownian motion. The scaling of the variance (i.e., how the mean squared displacement $\langle X^2 \rangle$ of the process X increases in time) is related to d_S .

To make sense of this picture in a spacetime context with D topological dimensions (one timelike and $D - 1$ spacelike), one must make a few strong assumptions on this diffusive process:

- (A) Replace the Laplacian ∇^2 with the covariant Laplace–Beltrami (or d’Alembertian) operator $\square = \nabla_\mu \nabla^\mu$ in imaginary time. Thus, there are D directions parametrized by coordinates x^μ , $\mu = 1, \dots, D$, where $x^D = it$ and $t = x^0$ is time in Lorentzian signature. In order to keep the probabilistic interpretation of the diffusion equation, the spatial generator is assumed to be an elliptic operator, hence the requirement of Euclideanization.
- (B) Time is thus treated on equal footing with the other coordinates, in conformity with the spirit of general relativity. Consequently, the variable t in the diffusion equation (1) is replaced by an abstract evolution parameter σ everywhere.
- (C) In particular, σ has the same dimensionality as t (it is a length or time scale, $[\sigma] = -1$ in momentum units), and the diffusion operator ∂_t in (1) is simply replaced by ∂_σ .

This construction may be unsatisfactory for various reasons. In general-relativistic systems, the diffusion equation is expected to have a covariant form, which should survive even under the simple replacement procedure (B)-(C). Yet, a non-relativistic diffusion equation is typically assumed for curved manifolds. Second, statistical-mechanics time t is Euclideanized in the spatial generator $\square$, but not in the diffusion operator (which would result in a Schrödinger equation). The general feeling is that, even when the diffusion equation is more or less well motivated in the context of statistical mechanics, it carries a considerable level of arbitrariness when promoted to the diffusion equation associated with a given spacetime. In this sense, the diffusion equation (and, hence, the spectral dimension d_S) of a spacetime cannot be truly derived from solid first principles.

For ordinary spacetimes, the proof of the pudding is in the eating, and the results of the procedure (A)-(C) are reasonable enough not to require a revision. We can better appreciate the problem, however, when moving to *multi-scale spacetimes*, in particular those with multi-fractional measure [7, 8, 9, 10, 11, 12, 13, 14, 15, 16]. These spacetimes [7, 17] have been recently introduced as realizations of geometries with anomalous (or even fractal-like) properties [18, 19, 20], which appear in certain regimes of many quantum-gravity models (a canonical list of references can be found, e.g., in [9, 13, 18, 21, 22]). The spectral dimension of multi-fractional spacetimes has been computed in [19, 8, 16] using a certain *Ansatz* for the diffusion equation, and found to be anomalous ($d_S \neq D$). Such diffusion equation, however, was simply assumed and it turns out to have some issues, which we discuss here. We completely revisit the problem (and address these issues) from the more fundamental point of view of statistical and quantum mechanics. Although the resulting diffusion equation will be different from the old assumption, the spectral dimension will still be anomalous, i.e., different from the integer topological dimension of spacetime. However, one must impose certain conditions in order for these anomalous spacetimes to be fractal in the usual sense. In practice, only one multi-scale model (with so-called q -Poincaré symmetries) will satisfy all these conditions.

As in [12, 13, 23], one of the key messages of the present work is that the spectral dimension is only one of the many possible ways to characterize a spacetime geometry: it is well known that an anomalous correlation function, whose exponent is governed by d_S , can be obtained by a number of inequivalent diffusion processes [24]. To remove this degeneracy, it is important to gain insight into the type of stochastic process underlying the diffusion equation. Thus, *spacetime geometries can be classified not only according to their Hausdorff and spectral dimension, but also by a stochastic process*. More generally, there exists a well-furbished ‘alternative toolbox’ [7, 13, 15] mutuanted from various disciplines of mathematics and mathematical physics (chaos theory, probability theory, transport and percolation theory, statistical mechanics, complex systems, multi-fractal geometry) which is still to be fully tapped into by the quantum gravity community, where discussions are often limited only to a few geometric indicators (Hausdorff and spectral dimension). We will provide a graphic representation of the possibilities of this arsenal by plotting the trajectories of random walkers associated with different fractional geometries.

The plan of the paper is as follows. In section 2 we review the case of ordinary Minkowski spacetime, starting from the derivation of the diffusion equation under the assumption of having a random walker with a certain stochastic interpretation (Brownian motion). This standard setting will introduce the main tools of statistical mechanics and probability and transport theory needed also in multi-scale spacetimes, the latter reviewed in section 3. In particular, in section 3.2 we analyze the diffusion equation previously assumed for these spacetimes and its related problems. Section 4 presents the new results for various classes of fractional spacetimes, using and extending the ‘alternative toolbox’ introduced in section 2. The spectral dimension of multi-scale geometries is computed in section 5, where it is shown that the theory with weighted Laplacian can have an ultraviolet regime $d_S \sim 0$ provided one interprets the scale in the measure as one signalling a transition between a ‘fuzzy’ and a continuous regime. Section 6 is devoted to discussion. Some of the results related to anomalous scaling can be obtained independently (and in agreement with the methods of the main text) from quantum mechanics; these digressions are contained in two appendices.

2. Brownian motion, diffusion equation and spectral dimension

The spectral dimension of a smooth classical manifold is an indicator of the geometry and topology of spacetime. It is obtained by letting a test particle diffuse on the Euclidean version of the manifold, and calculating the probability to find the probe again at the starting point after some diffusion time σ (not to be confused with coordinate time t , which is treated on the same ground as a spatial coordinate). This return probability is a function $Z(\sigma)$ of diffusion time; if $Z \sim \sigma^{-d_S/2}$ is a power law, the spectral dimension is the exponent d_S .

The test particle is assumed to follow a random walk of Brownian type. This is because the associated probability density function yields, in the absence of curvature,

the correct spectral dimension $d_S = D$, coinciding with the topological dimension of spacetime. The diffusion equation of Brownian motion [25, 26, 27] (see [28, 29] for extensive presentations) can be derived from a Langevin equation. We review Brownian motion and its diffusion equation in a non-relativistic space in sections 2.1 and 2.2. Promotion to the spacetime picture according to (A)-(C) will take place in section 2.3.

2.1. Brownian motion

Let $X(t)$ be the random variable denoting the position of a particle at time t ; in D dimensions, it is a vector. A Wiener process (or Brownian motion) $X_{\text{BM}}(t)$ with initial condition $X_{\text{BM}}(0) = x'$ is such that:

- (i) X_{BM} is continuous in t almost surely (i.e., with probability 1);
- (ii) the increments of X_{BM} are uncorrelated, meaning that $X_{\text{BM}}(t_2) - X_{\text{BM}}(t_1)$ is independent of $X_{\text{BM}}(t_4) - X_{\text{BM}}(t_3)$ if the intervals (t_1, t_2) and (t_3, t_4) do not overlap, $(t_1, t_2) \cap (t_3, t_4) = \emptyset$;
- (iii) X_{BM} is governed by a Gaussian distribution.

To make the last requirement explicit, let us introduce a dimensionless white noise $\eta(t)$, that is, a Gaussian random field such that

$$\langle \eta(t) \rangle_\eta = 0, \quad \langle \eta(t) \eta(t') \rangle_\eta = \kappa_1 \delta(t - t'), \quad (2)$$

where κ_1 is a constant with engineering dimension $[\kappa_1] = -1$. The noise is called ‘white’ because its spectrum (the Fourier transform of its two-point correlation function) is a constant independent of the frequency. Here, angular brackets with subscript η denote the average over the stochastic background, i.e., with respect to the PDF $u(\eta, t)$ of the process: for any f , $\langle f(\eta, t) \rangle_\eta := \int d^D \eta u(\eta, t) f(\eta, t)$. In the absence of external forces, the *Langevin equation* of Brownian motion is

$$\partial_t X_{\text{BM}}(t) = \eta(t). \quad (3)$$

By definition, it represents a Wiener process as the integral of a white noise:

$$X_{\text{BM}}(t) := x' + \int_0^t dt' \eta(t'). \quad (4)$$

This way, the differential dX_{BM} is well defined even if the trajectory $X_{\text{BM}}(t)$ is nowhere differentiable in the ordinary sense. Two key features of Brownian motion are that it obeys the scaling property

$$X_{\text{BM}}(\lambda t) = \lambda^{\frac{1}{2}} X_{\text{BM}}(t), \quad (5)$$

and that it possesses stationary increments, i.e.,

$$\langle [X(t) - X(t')]^2 \rangle = \langle X^2(t - t') \rangle. \quad (6)$$

Here, angular brackets without subscript denote the average with respect to the PDF $\hat{P}$ of the process, $\langle f(X, t) \rangle := \int d^D X \hat{P}(X, x', t) f(X, t)$.

2.2. Diffusion equation

A Fokker–Planck equation (namely, the diffusion equation) for the PDF of Brownian motion can be derived starting from the Langevin equation [28, 29]

$$m\ddot{X} + m\gamma\dot{X} + U'(X) = F, \quad (7)$$

where $[m] = 1 = [\gamma]$, $[F] = 2$, and dots and primes denote derivatives with respect to, respectively, time t and X . This equation describes the motion of a particle $X(t)$ of mass m in the presence of a friction force (proportional to the mass of the particle, a constant coefficient γ and the velocity $V = \dot{X}$), an external potential U , and a random force F , representing the pushing around of the particle by the medium. A writing of equation (7) where the left- and right-hand sides are dimensionless is

$$\frac{1}{\gamma}\ddot{X} + \dot{X} + \frac{U'(X)}{m\gamma} = \eta, \quad \eta := \frac{F}{m\gamma}. \quad (8)$$

There are various methods to find the diffusion equation. One takes the Langevin equation (8) exactly and solves it. Consider for instance the case without potential, $U = 0$. Integrating twice the Langevin equation $\dot{V} + \gamma V = \gamma\eta$ starting from time $t = 0$, one obtains

$$\begin{aligned} X(t) &= \int_0^t dt' V(t') \\ &= x' + V_0\chi(t) + \gamma \int_0^t dt'' \chi(t-t'')\eta(t''), \end{aligned} \quad (9)$$

where $V_0 = V(0)$ is the initial velocity and

$$\chi(t-t'') := \frac{1 - e^{-\gamma(t-t'')}}{\gamma}. \quad (10)$$

Expression (9) holds true also in more general situations with a more complicated function χ , for instance when $U \neq 0$, when the friction term $m\gamma\dot{X}$ in (7) is replaced by a non-local operator and the process is non-Markovian, and for general colored noise η . If the stochastic properties of η are known, the PDF's resulting from these generalized Langevin equations (GLE) [29, 30, 31, 32, 33, 34, 35, 36, 37] can be found by elegant techniques, for instance using the characteristic (or generating) functional approach [28, 31, 33], or the functional derivative approach [32], or the conservation-law approach [29]. In particular, the diffusion equation for Brownian motion (also known as the Smoluchowski equation when $U = 0$) is obtained from equation (9) under three assumptions: (i) that η is a Gaussian white noise, equation (2) (thus, also X and the velocity field V are Gaussian); (ii) in the limit $V_0 \rightarrow 0$ (Maxwell distribution of velocities); (iii) assuming to look at times much larger than the relaxation time $t_{\text{rt}} = 1/\gamma$. The latter corresponds to taking $\chi \approx \text{const}$ from the start. Namely, setting $U = 0$ and dropping the second-derivative term in (8), we have $\dot{X} \approx \eta$, which is nothing but (3).

In fact, for our purposes it suffices to apply the conservation-law approach [29], valid for Markovian systems, and to make direct use of equation (3). Although the following steps are well known, we report them to illustrate the relation between Langevin and

diffusion equation for later use. The starting point is the conservation of the probability density,

$$\langle 1 \rangle = \int d^D X \hat{P}(X, x', t) = 1 \quad \forall t. \quad (11)$$

Differentiating this expression with respect to t implies that $\partial_t \hat{P}$ is proportional to a total divergence, in particular (as it happens in fluid mechanics, statistical mechanics or electrodynamics) to the divergence of its flux $\dot{X} \hat{P}$. Thus, the conservation law is $\partial_t \hat{P} + c \nabla_X \cdot (\dot{X} \hat{P}) = 0$, for some constant c . From equation (3), we can rewrite this expression as

$$\partial_t \hat{P} + c \nabla_X \cdot (\eta \hat{P}) = 0. \quad (12)$$

Integrating in time, one has

$$\hat{P}(X, x', t) = P_0(X, x') - c \int_0^t dt' \nabla_X \cdot [\eta(t') \hat{P}(X, x', t')], \quad (13)$$

where $P_0(X, x') = P(X, x', 0)$ is the initial condition. For times prior to t , the functional $\hat{P}$ depends implicitly on the noise η . Plugging equation (13) back into (12), we find

$$\partial_t \hat{P} = -c \nabla_X \cdot (\eta P_0) + c^2 \nabla_X \cdot \left[\eta \int_0^t dt' \nabla_X \cdot (\eta \hat{P}) \right] = 0. \quad (14)$$

We take the average over the noise and call P the stochastic average of the PDF (which depends on η , via equation (12)):

$$P(x, x', t) := \langle \hat{P} \rangle_\eta. \quad (15)$$

The Gaussian statistics of the white noise now leads to the desired result. In fact, equation (2) implies that the average of the first term in the right-hand side of (14) vanishes (X and η are independent variables), while the bracket in the second term of (14) is made of two contributions:

$$\int_0^t dt' \nabla_X \cdot [\langle \eta(t) \eta(t') \rangle_\eta P(x, x', t')] + \int_0^t dt'' [\eta(t'') \langle \eta(t) \nabla_X \cdot \hat{P}(X, x', t'') \rangle_\eta], \quad (16)$$

since $\hat{P}$ does include noise factors at time t' . The first correlation function gives a $\delta(t - t')$, the second a $\delta(t - t'')$ for $t'' < t'$. But $t > t''$ strictly, so the second contribution vanishes and, setting $c = 1$ without loss of generality, we are left with

$$(\partial_t - \kappa_1 \nabla_x^2) P(x, x', t) = 0, \quad (17)$$

where we have thrown away higher-order noise terms so that the Laplacian acts on the spatial coordinates x .

The diffusion coefficient κ_1 is measured in units of $(\text{length})^2/(\text{time})$. Often it is effectively absorbed in the definition of a length variable

$$\ell(t) := \sqrt{\kappa_1 t}. \quad (18)$$

In general, the diffusion coefficient is determined by the stochastic process underlying the diffusion equation (e.g., [28, 38, 39]). Having averaged over noise, the average $\langle \cdot \rangle$ with

respect to $\hat{P}$ and the one with respect to P coincide, $\langle f(x, t) \rangle := \int d^D x P(x, x', t) f(x, t)$, and they shall be denoted in the same way without subscript. In particular, one can show that the mean squared displacement (or second moment, or variance) from $x' = 0$ is, in D dimensions,

$$\langle X_{\text{BM}}^2(t) \rangle = 2D\kappa_1 t. \quad (19)$$

The Laplacian ∇^2 in equation (17) acts on the x dependence of the solution P , but in this particular case the latter depends only on the distance $r = |x - x'|$ between the initial and final point. In ordinary Euclidean space, the solution is the Gaussian distribution

$$P(x, x', t) = u_1(x, x', t) := \frac{e^{-\frac{|x-x'|^2}{4\ell^2(t)}}}{[4\pi\ell^2(t)]^{\frac{D}{2}}}, \quad (20)$$

where $|x - x'|^2 = (x_1 - x'_1)^2 + \dots + (x_D - x'_D)^2$ and the squared length ℓ^2 is the dispersion of the Gaussian. Consistently with (11), the solution is a probability density with normalization

$$\int_{-\infty}^{+\infty} d^D x u_1(x, x', t) = 1, \quad (21)$$

meaning that at all times or scales the pointwise test particle (represented by the delta initial condition $u_1(x, x', 0) = \delta(x - x')$) can be almost surely found somewhere.

2.3. Spectral and walk dimension

So far we have interpreted the variable t , which has dimension $[t] = -1$, as a time parameter. Moving away from transport theory, in the context of spacetime theories $t \rightarrow \sigma$ is simply a length, representing the characteristic length scale ℓ (equation (18) with $t \rightarrow \sigma$) at which one is probing the geometry. The Laplacian is then replaced by the curved Laplace–Beltrami operator in Wick-rotated spacetime.

When a stochastic process is staged on a smooth manifold with boundary, the return probability (or heat kernel, or partition function) $Z(\sigma)$ is simply defined as the functional trace of the solution P , i.e., its integral over the volume of the set when $x = x'$. Then, the first term in the Seeley–DeWitt expansion (e.g., [40, 41, 42]) is of the form $Z(\sigma) = \mathcal{V}/(4\pi\kappa_1\sigma)^{-d_s/2} + \dots$, where $\mathcal{V}$ is the volume of the set. In the present case, however, this volume is infinite and it is customary to define the return probability as the trace per unit volume:

$$\mathcal{P}(\sigma) = \frac{Z(\sigma)}{\mathcal{V}} = \frac{\int d^D x P(x, x, \sigma)}{\int d^D x}. \quad (22)$$

This definition coincides with the spatial average of the return probability density $P(x, x, \sigma)$. Since $P(x, x, \sigma) = u_1(x, x, \sigma)$ is constant in x for a Brownian motion, the denominator in equation (22) exactly cancels the divergence in the numerator and one ends up with $\mathcal{P}(\sigma) = [4\pi\ell^2(\sigma)]^{-D/2} \propto \sigma^{-D/2}$. The spectral dimension is simply $d_s = D$ and it determines the decaying law of the return probability. In general, it is defined as

$$d_s(\sigma) = -2 \frac{d \ln \mathcal{P}(\sigma)}{d \ln \sigma}, \quad (23)$$

or, more formally (due to a hidden divergence),

$$d_S(\sigma) := -2 \frac{d \ln Z(\sigma)}{d \ln \sigma}. \quad (24)$$

Equation (24) is equivalent to the alternative definition $\tilde{d}_S := -d \ln Z(\ell)/d \ln \ell$ only for normal diffusion, but in general the correct one is (24) [1, 2, 3, 4, 5, 6]. Dimensionally, the mean squared displacement $\langle X^2(\sigma) \rangle$ is always proportional to some squared length $\ell^2(\sigma)$.

These definitions are meaningful in a spacetime geometry context provided σ (or ℓ) is interpreted as a measured length scale. Then, d_S is the scaling law of the return probability when a test particle is let diffuse on the manifold. In this respect, one should notice that curvature and topology effects do modify the value $d_S = D$ at scales σ larger than the characteristic curvature radius. The classic example is the sphere: at $\sigma \sim 0$, the probe locally feels a Euclidean plane, but as time passes the compact topology ‘helps’ the particle going back to the initial point, so that $d_S \sim 0$ in the limit $\sigma \rightarrow +\infty$. Therefore, when the spectral dimension of spacetime is computed, the tacit understanding is that one is looking for the local geometric properties of the manifold, so that in all the cases where Z is not expected to be a simple power law due to topology or curvature, it is more convenient to define

$$d_S := -2 \lim_{\sigma \rightarrow 0^+} \frac{\ln Z(\sigma)}{\ln \sigma}. \quad (25)$$

Clearly, when Z is a power law this expression coincides with (24) and d_S is constant.

In more exotic scenarios of quantum gravity the effective diffusion equation can be very complicated [13], but at least in semi-classical regimes it must admit a positive semi-definite solution P . Otherwise, the operational definition of the spectral dimension (24) as the scaling of the return probability associated with a random walk would be lost. The requirement $P \geq 0$ was advocated, in particular, in [13, 23].

The variance defines the so-called *walk dimension*:

$$\langle X^2 \rangle \propto \sigma^{2/d_W}. \quad (26)$$

From equation (19), it follows that $d_W = 2$ (normal diffusion) for ordinary Minkowski spacetime. In general, the spectral dimension d_S precisely determines the anomalous scaling in σ , since there is a relation between Hausdorff, spectral, and walk dimension [6]:

$$d_W = 2 \frac{d_H}{d_S}. \quad (27)$$

Consistently, for Minkowski spacetime $d_H = D = d_S$. This relation holds for all fractals and can be understood from (26) in various ways [6]. A particularly simple one is to consider two random-walk systems characterized, respectively, by a length ℓ (the root-mean-square displacement $\ell \sim \sqrt{\langle X^2 \rangle}$ of the walker) and a rescaled one $\ell \rightarrow \lambda \ell$. The energy density of states $\rho_\ell(E)$ is an extensive quantity proportional to the Hausdorff volume of the system, $\rho_\ell(E) \sim \ell^{d_H}$. Therefore, under rescaling $\ell \rightarrow \lambda \ell$ one has

$$\rho_{\lambda \ell}(E) = \lambda^{d_H} \rho_\ell(E). \quad (28)$$

On the other hand, from the Schrödinger equation one sees that energy E is dimensionally conjugate to time, but from equation (26) ($\sigma \sim \ell^{d_W}$) there follows that

$$E_{\lambda\ell} = \lambda^{-d_W} E_\ell. \quad (29)$$

For probability to be conserved under rescaling, the density of states in the original and rescaled system must be related to each other by $\rho_{\lambda\ell}(E_{\lambda\ell})dE_{\lambda\ell} = \rho_\ell(E_\ell)dE_\ell$, which implies, from equation (29),

$$\rho_{\lambda\ell}(E_{\lambda\ell}) = \lambda^{d_W} \rho_\ell(E_\ell). \quad (30)$$

The solution of equations (28)–(30) is (removing the subscript ℓ from now on)

$$\rho(E) \sim E^{d_H/d_W-1} =: E^{d_S/2-1}, \quad (31)$$

where in the last step we defined the spectral dimension in analogy with the ordinary Euclidean-space expression $\rho(E) \sim E^{D/2-1}$. Via quantum mechanics, one can then show (not so straightforwardly) that the definition in (31) of d_S coincides with the one from the diffusion equation.

3. Multi-scale spacetimes

A generic spacetime $\mathcal{M}$ with scale-dependent geometry [18, 7] can be defined by equipping ordinary Minkowski spacetime with a measure $d\varrho(x)$, replacing the ordinary Lebesgue measure $d^D x$. A non-trivial metric structure independent of the measure can be added to describe manifolds with curvature, but we shall ignore it here, as we are interested in how a change of the differential structure (and of momentum space) affects the properties of spacetime. For technical reasons [14, 16], it is convenient to concentrate on *factorizable* measures,

$$d\varrho(x) := d^D x v(x) = dt v_0(t) d\mathbf{x} v(\mathbf{x}) := dt v_0(t) \prod_{i=1}^{D-1} dx^i v_i(x^i), \quad (32)$$

where the D functions v_μ can be all different. By definition, coordinates have dimension of lengths ($[x] = -1$ in momentum units). A further assumption is that the measure weight be positive semi-definite, $v_\mu = v_\mu(x^\mu) \geq 0$ for all $\mu = 0, 1, \dots, D-1$. An example of factorizable measure is the fractional one,

$$d\varrho_\alpha(x) = d^D x v_\alpha(x), \quad v_\alpha(x) = \prod_\mu v_\alpha(x^\mu) := \prod_\mu \frac{|x^\mu|^{\alpha_\mu-1}}{\Gamma(\alpha_\mu)}, \quad (33)$$

where Γ is the gamma function and α_μ are D real parameters (‘fractional charges’) in the range $0 < \alpha_\mu \leq 1$. In the simplest ‘isotropic’ case, $\alpha_\mu = \alpha$ for all μ . In general, we will call α the average fractional charge

$$\alpha := \frac{1}{D} \sum_\mu \alpha_\mu. \quad (34)$$

Fractional measures are not unique (the position of the singularity parametrizes an infinite class, there exist non-factorizable versions as in the Riesz integral, and so on [8])

but they epitomize the simplest class of continuous measures with anomalous scaling law $\varrho_\alpha(\lambda x) = \lambda^{d_H} \varrho_\alpha(x)$, where d_H is the Hausdorff dimension of spacetime. For equation (33), for instance,

$$d_H = \sum_{\mu} \alpha_{\mu} = D\alpha \leq D. \quad (35)$$

To obtain a scale-dependent geometry, it is sufficient to sum over a finite set of charges α [7, 9]. Keeping factorizability, the multi-scale measure weight is [16]

$$v_*(x) := \prod_{\mu} v_*(x^{\mu}) := \prod_{\mu} \left[\sum_n g_n v_{\alpha_n}(x^{\mu}) \right], \quad (36)$$

where $g_n \geq 0$ are dimensionful coefficients depending on a hierarchy of length scales $\ell_n > 0$. For example, an isotropic spacetime with Hausdorff dimension $d_H \sim D$ in the infrared (IR) and $d_H \sim D\alpha_* < D$ in the ultraviolet (UV) is characterized by two charges $0 < \alpha_1 = \alpha_* < 1$ and $\alpha_2 = 1$ and one fundamental length $\ell_1 = \ell_*$ defining large and small scales. The measure weight then reads, for each coordinate,

$$v_*(x) = 1 + \ell_*^{1-\alpha_*} v_{\alpha_*}(x). \quad (37)$$

The volume of a D -ball of radius R can be easily calculated and is of the form $\mathcal{V}^{(D)} \sim \ell_*^D [\Omega_{D,1}(R/\ell_*)^D + \Omega_{D,\alpha_*}(R/\ell_*)^{D\alpha_*}]$, where $\Omega_{D,\alpha}$ is the fractional volume of a unit ball ($\Omega_{D,\alpha} = \Omega_{D,1}/[\Gamma(\alpha_* + 1)]^D$ if centered at the origin) and we have thrown away off-diagonal terms in the integration. When $R \gg \ell_*$, the first term dominates and the ball volume scales as usual; otherwise, the second term dominates and the UV scaling is anomalous. In particular, if $\alpha_* = 2/D$ spacetime has $d_H = 2$ in the UV.

A more fundamental version of fractional spacetimes is endowed with complex fractional measures [7, 9, 11]. By a suitable choice of coefficients, one can construct measures with log-oscillations, of which the present real-order measures are nothing but the zero mode. Log-oscillating measures describe spacetimes with discrete symmetries and a hierarchy of ultra-microscopic characteristic scales, smaller than those appearing in the real-order multi-fractional measure (36). At the bottom of this tower, there is a fundamental length possibly identified with Planck's length [11]. The interest in these models lies in the phenomenological applications of such a scale hierarchy in quantum gravity, as well as on the fact that complex fractional measures better represent deterministic fractals [43]. When defining the spectral dimension for these geometries, it is necessary to average over the log-period of the measure in order to get a meaningful observable [9]. The only surviving contribution is the zero mode, i.e., a power-law with real fractional charge. Therefore, all information about the spectral dimension is encoded in the models considered in the present work, and we do not need to include log-oscillations in the discussion.

3.1. Laplacians

We consider three classes of multi-fractional spacetimes, each characterized by a different symmetry for the Lagrangian density and, hence, by a different Laplace–Beltrami

(Laplacian in short) operator. In the remainder of this section, we leave the form of the weight $v(x)$ arbitrary while assuming the factorizability and positivity conditions.

3.1.1. Weighted Poincaré symmetries With respect to the natural scalar product on $\mathcal{M}$, one can construct a class of self-adjoint operators $\mathcal{K}_{v,\gamma}$ of order 2γ generalizing the d'Alembertian [13]. In particular, the second-order operator belonging to this class is [10]

$$\mathcal{K}_v := \eta^{\mu\nu} \mathcal{D}_\mu \mathcal{D}_\nu, \quad \mathcal{D}_\mu := \frac{1}{\sqrt{v(x)}} \partial_\mu \left[\sqrt{v(x)} \cdot \right], \quad (38)$$

where $\eta_{\mu\nu} = \text{diag}(-, +, \dots, +)$ is the Minkowski metric. A field action endowed with this operator is invariant under deformed Poincaré symmetries, where the generator algebra is the usual one in the free case, but the algebra elements do not generate the standard Poincaré transformations [16]. In the interacting case, the algebra itself is deformed.

The eigenfunctions of $\mathcal{K}_v$ are (here $k^2 := k_\mu k^\mu = -k_0^2 + \sum_{i=1}^{D-1} k_i^2$)

$$\mathfrak{e}(k, x) = \frac{1}{\sqrt{w(k)v(x)}} \frac{e^{ik \cdot x}}{(2\pi)^{\frac{D}{2}}}, \quad \mathcal{K}_v \mathfrak{e}(k, x) = -k^2 \mathfrak{e}(k, x), \quad (39)$$

where w is the measure weight of momentum space. The normalization is chosen so that one can write an invertible momentum transform:

$$\tilde{f}(k) := \int_{-\infty}^{+\infty} d^D x v(x) f(x) \mathfrak{e}^*(k, x), \quad (40)$$

$$f(x) = \int_{-\infty}^{+\infty} d^D k w(k) \tilde{f}(k) \mathfrak{e}(k, x). \quad (41)$$

The multi-scale generalization of the Dirac distribution is then

$$\delta_v(x, x') := \frac{\delta(x - x')}{\sqrt{v(x)v(x')}} = \int d^D k w(k) \mathfrak{e}^*(k, x) \mathfrak{e}(k, x'), \quad (42)$$

with a similar expression for $\delta_w(k, k')$. The operator (38) was considered in [8, 9, 10, 12, 13, 14, 15, 16].

3.1.2. Ordinary Poincaré symmetries In another scenario [18, 19, 20, 8, 9] the Lagrangian density possesses ordinary Poincaré symmetries and the Laplace–Beltrami operator is simply

$$\square := \eta^{\mu\nu} \partial_\mu \partial_\nu, \quad (43)$$

which is not self-adjoint with respect to the scalar product with measure weight $v(x)$. In fact,

$$\square^\dagger = \check{\mathcal{K}}_v := \frac{1}{v(x)} \square [v(x) \cdot]. \quad (44)$$

Hence, a (for instance) scalar field model with quadratic-like kinetic term $-\partial_\mu \phi \partial^\mu \phi / 2$ is inequivalent to the one with Gaussian-like term $\phi \square \phi / 2$. In this case, the field theory has

some ordering prescription and there is no direct definition of a self-adjoint momentum operator (and there may appear complications with the micro-causality structure of the theory [9]). Correspondingly, it is not clear whether an invertible momentum transform exists. Still, the operators (43) and (44) will be of interest in what follows.

3.1.3. q -Poincaré symmetries The action measure can be recast as $d\varrho = d^D q$, where $dq^\mu = dx^\mu v_\mu(x^\mu)$. Notice that this writing, as well as the most general one $q(x) = \varrho(x)$, is equivalent to equation (33) only in the sense of distributions. In the case of a fractional spacetime with fixed dimensionality, for each direction one has

$$q^\mu(x^\mu) = \varrho_\alpha(x^\mu) = \frac{\text{sgn}(x^\mu)|x^\mu|^{\alpha_\mu}}{\Gamma(\alpha_\mu + 1)}. \quad (45)$$

Obviously, the Hausdorff dimension is not affected by a change of variables [8], and it is equation (35), $d_H = D\alpha$. This happens because *by definition* the momentum space of the theory is conjugate to position space in x , not in q . Consequently, the units of the x coordinates are $[x^\mu] = -1$ while the q 's have anomalous scaling, $[q^\mu] = -\alpha_\mu$. This feature guarantees that the theory is not trivial [15] even if one defines it such that in q variables it is formally identical to the ordinary one, including the Laplace–Beltrami operator

$$\square_{q(x)} = \eta^{\mu\nu} \frac{\partial}{\partial q^\mu(x)} \frac{\partial}{\partial q^\nu(x)} = \eta^{\mu\nu} \frac{1}{v_\mu(x)} \partial_\mu \left[\frac{1}{v_\nu(x)} \partial_\nu \cdot \right], \quad (46)$$

which is, therefore, self-adjoint under suitable boundary conditions. In the last equation we omitted the indices in the arguments of q and v .

The measure, the Lagrangian, and the action as a whole possess ‘ q -Poincaré’ symmetries, i.e., they are Lorentz and translation invariant under transformations over the q 's, which are then non-linear transformations over the x 's [8, 9]:

$$q'^\mu(x') = \Lambda^\mu_\nu q^\nu(x) + a^\mu. \quad (47)$$

3.2. Previous Ansatz for the diffusion equation

The explicit calculation of the spectral dimension d_S was performed in [8] for fixed dimensionality and in [9, 13, 16] for multi-fractional spacetimes, in the case of the weighted Laplacian (38).[‡] We recall here the result for an integer-order diffusion equation and an isotropic fractional measure ($\alpha_\mu = \alpha$, fixed dimensionality). The multi-scale anisotropic case is more complicated and adds nothing to the main point we wish to make here.

The Ansatz adopted for the diffusion equation was based on two assumptions: (a) since σ is a fictitious variable, the diffusion operator ∂_σ should not reflect the differential structure of fractional spacetime and can be left unchanged; (b) as the spatial generator in the ordinary diffusion equation (17) coincides with the self-adjoint

[‡] A heuristic estimate of d_S for the ordinary Laplacian can be found in [19].

Laplacian appearing in the action, so should it be in the fractional case. This singles out the diffusion equation

$$(\partial_\sigma - \kappa_1 \mathcal{K}_v) \tilde{P}(x, x', \sigma) = 0, \quad \tilde{P}(x, x', 0) = \delta_v(x, x'). \quad (48)$$

The first problem arises because the solution $\tilde{P}(x, x', \sigma) = u_1(x, x', \sigma) / \sqrt{v(x)v(x')}$ is not normalized to 1. It is easy to see that $P := \sqrt{v(x')/v(x)} \tilde{P} = u_1/v(x)$ obeys the same initial condition, is normalized to 1, and is solution of

$$(\partial_\sigma - \kappa_1 \check{\mathcal{K}}_v) P(x, x', \sigma) = 0, \quad (49)$$

rather than of equation (48) [13]. The operator $\check{\mathcal{K}}_v$ is not self-adjoint, which requires to abandon assumption (b). This is not an issue, since Fokker–Planck equations are in general not self-adjoint [29]. One should now explain, however, why the spatial generator carries weights v instead of $\sqrt{v}$, but for the time being we just ignore this point and move on.

In analogy with equation (22), we maintain the definition of return probability as the functional trace of the solution P per unit volume:

$$\mathcal{P}(\ell) = \frac{Z(\ell)}{\mathcal{V}_H} = \frac{1}{\mathcal{V}_H} \int d\varrho(x) P(x, x, \ell), \quad \mathcal{V}_H := \int d\varrho(x) = \int d^D x v(x). \quad (50)$$

In fractional spacetimes, however, the spacetime integral in the denominator does not cancel the one in the numerator. In fact, the return probability reads

$$\mathcal{P}(\ell) = \frac{\int d^D x}{\int d^D x v(x)} \frac{1}{(4\pi\ell^2)^{\frac{D}{2}}}. \quad (51)$$

The ratio of the two spatial integrals is a divergent, dimensionful constant. To get rid of it, one must first extract all the dimensional dependence. This is done by defining the dimensionless coordinates $\tilde{x}^\mu = x^\mu/\ell$, so that $\mathcal{P}(\ell) = C\ell^{-D\alpha}$. The constant C is formally divergent but it can be regularized [13] and, since it is dimensionless, be thrown away. It is completely immaterial in the definition (23), which would give

$$d_S = D\alpha. \quad (52)$$

Unfortunately, there is an issue with this procedure, which we can state in several ways. One is to notice that the regularization trick would introduce a degree of arbitrariness in the return probability which, if the underlying stochastic process is well defined, should not be there. In other words, the definitions (23) and (24) are inequivalent in this case, while they should always agree because the overall volume prefactor in $\mathcal{P}$ does not depend on diffusion time. This would suggest to use (24) or, which is the same, to define $\mathcal{P}$ as $\mathcal{P} = Z/(\int d^D x)$ instead of equation (50), thus modifying the result of the spectral dimension to $d_S = D$. This is not even a fractal, since $d_S = D \geq d_H$ (for fractals, $d_S \leq d_H$). The interpretation of $\mathcal{P}$ as the spatial average of P would also be lost. On the other hand, one might waive away worries that the overall constant in the return probability (51) is regularization-dependent on account of two observations. First, in scenarios with fixed dimensionality the regularization only affects the normalization of the PDF, not its signature. Second, when d_S is multi-scale,

transient regimes between asymptotic plateaux may also depend on the regularization, which is a known feature of multi-scale systems including field theories of quantum gravity [13].

Another way to see the problem is to recall that the heat kernel for fractals is of the form

$$Z(\ell) = \frac{\mathcal{V}_H}{(4\pi\ell^2)^{d_S/2}} + \dots, \quad (53)$$

where $\mathcal{V}_H = L^{d_H}$ is the spectral volume of the set, L is the characteristic spectral length determined by the Laplacian (roughly, by the periodicity of its eigenfunctions) and d_H is the Hausdorff dimension of the set [44, 45]. On the other hand, the divergent contribution from the heat kernel Z in (51) is the ordinary integer volume, not the Hausdorff volume. This version of the problem can be mildened by recalling that the first Seeley–DeWitt coefficient in the heat kernel is quite generically equal to the spectral volume but, to the best of our knowledge, it has not been proven as a strict rule for non-smooth sets. It is not obvious, however, why this should not be the case also for fractional spacetimes, which are a hybrid between continuum geometries and genuine fractals.

A third manifestation of the problem becomes apparent when one identifies its origin in the normalization of plane waves (39): the phases $\mathfrak{e}(k, x)$ are normalized per unit integer volume, not per fractional volume. This is apparent from equation (42). In yet other words, the number of states is not the fractional volume in equation (50) but, rather, $\delta(0)/v(x)$. In fact, define the density of states of a system with Hamiltonian eigenstates $\psi_k(x)$ and energy E_k as

$$w(E) \rho(E) := \int d^D k w(k) |\psi_k(x)|^2 \delta(E - E_k), \quad (54)$$

where $w(E)$ is the energy measure weight. Then, the total number of states is

$$\int dE w(E) \rho(E) = \int d^D k w(k) |\psi_k(x)|^2 = \frac{\delta(0)}{v(x)}, \quad (55)$$

where in the last step we used the fact that the states ψ_k always contain a factor of the form $1/\sqrt{v(x)w(k)}$ (as in the free particle case, where $\psi_k(x) = \mathfrak{e}(k, x)$). In the bra-ket formalism, this is tantamount to starting with the resolutions of the identity

$$\mathbb{1}_k = \frac{1}{(2\pi)^{D/2}} \int d^D k w(k) |k\rangle \langle k|, \quad \mathbb{1}_x = \frac{1}{(2\pi)^{D/2}} \int d^D x v(x) |x\rangle \langle x|, \quad (56)$$

so that $|x\rangle = (2\pi)^{-D/2} \int d^D x' v(x') |x'\rangle \langle x'|x\rangle$ and one finds $\langle x'|x\rangle = (2\pi)^{D/2} \delta_v(x, x')$. Since $(2\pi)^{D/2} \delta_v(x, x') = \langle x' | \mathbb{1}_k | x \rangle$, the right-hand side of (55) can be recast as $\langle x | x \rangle = \delta(0)/v(x)$, where $\delta(0)$ is the volume of integer momentum space. It is the integer volume because the eigenfunctions $\psi_k(x)$ always cancel the weight factor $w(k)$. Therefore, each infinitesimal hypercube $d^D k$ (or $d^D x$) contains $w(k)d^D k/w(k) = d^D k$ localized states (respectively, $d^D x$), as in the usual case.

Ultimately, the problem lies in the diffusion equation itself, equation (49), which was simply assumed. Starting from the Langevin equation we will get, instead, a

different diffusion equation not only derived in closer conformity with the standard case of ordinary space, but also capable of overcoming the difficulties outlined above.

4. Diffusion in fractional spacetimes

4.1. Weighted Laplacian

The natural extension of classical mechanics to multi-scale spacetimes is obtained by replacing ordinary time derivatives with weighed derivatives $\mathcal{D}_t$ [14]. For instance, a free particle in a quadratic potential has action

$$S = \int dt v_0(t) \left[\frac{1}{2} m (\mathcal{D}_t x)^2 - \frac{1}{2} m \omega^2 x^2 \right], \quad (57)$$

with equation of motion

$$m \mathcal{D}_t^2 x + m \omega^2 x = 0. \quad (58)$$

Similarly, it is natural to define the Langevin equation for a random variable X in fractional or multi-scale space as

$$m \mathcal{D}_t^2 X + m \gamma \mathcal{D}_t X + U'(X) = F. \quad (59)$$

Switching to the spacetime interpretation (A)-(C) spelled in the introduction, one ends up with the stochastic equation

$$m \mathcal{D}_\sigma^2 X + m \gamma \mathcal{D}_\sigma X + U'(X) = F, \quad (60)$$

where we switched to diffusion time according to assumption (B). However, now assumption (C) is less transparent than in the standard case, since the derivative $\mathcal{D}_\sigma$ contains a weight $v(\sigma)$ for an abstract evolution parameter and, *a priori*, it is not obvious whether this weight should have the same functional form as the weight $v_0(t)$. For the time being, we leave the form of $v(\sigma)$ unspecified, calling $1 - \beta = [v(\sigma)]$ its scaling dimension. In particular, in the fractional case we set

$$v(\sigma) = v_\beta(\sigma) := \frac{\sigma^{\beta-1}}{\Gamma(\beta)}, \quad \beta > 0, \quad (61)$$

with β not necessarily equal to the fractional charge α_0 .

Making now the field redefinition

$$Y(\sigma) := \sqrt{v(\sigma)} X(\sigma), \quad \xi(\sigma) := \frac{1}{m\gamma} \sqrt{v(\sigma)} F(\sigma), \quad (62)$$

equation (60) reduces to (8) in the presence of a non-autonomous potential $W = \sqrt{v(\sigma)} U$ and a noise ξ :

$$\frac{1}{\gamma} \partial_\sigma^2 Y + \partial_\sigma Y + \frac{W'(\sigma, Y)}{m\gamma} = \xi. \quad (63)$$

The resemblance with equation (8) is only superficial, since measure units differ: the effective variable Y has scaling dimension $[Y] = -(1 + \beta)/2$, while $[\xi] = (1 - \beta)/2$. Notice that if X is a random variable, so will Y , since $v(\sigma)$ is a deterministic function.

Also, at the initial time $\sigma = 0$ one takes the expression $Y(0) = \lim_{\sigma \rightarrow 0} \sqrt{v(\sigma)} X(\sigma)$ formally finite, even if $v(0)$ may diverge.

A natural assumption is that F is a Gaussian white-noise field but in a fractional sense, i.e., with correlation $\langle F(\sigma)F(\sigma') \rangle_F \propto \delta_v(\sigma, \sigma')$, where the proportionality coefficient has dimension $(\text{mass})^{4-\beta}$ ($\langle FF \rangle$ has scaling dimension 4 and $[\delta_v] = \beta$). It is more instructive, however, to consider the general case

$$\langle F(\sigma)F(\sigma') \rangle_F = (m\gamma)^2 \kappa_{\beta,\nu} \sigma^{\nu-1} \delta_v(\sigma, \sigma') \quad \Rightarrow \quad \langle \xi(\sigma)\xi(\sigma') \rangle_\xi = \kappa_{\beta,\nu} \sigma^{\nu-1} \delta(\sigma - \sigma') \quad (64)$$

for all σ and σ' , where ν is a real parameter and $\kappa_{\nu,\beta}$ is a constant with dimension $[\kappa_{\beta,\nu}] = \nu - \beta - 1$. Repeating the same calculation as in section 2, one finds an expression for the probability density $\hat{u}$ (we reserve the symbol $\hat{P}$ for later) satisfying an equation of the form (14), with X replaced by Y . Taking the average over the noise and calling $u(y, y', \sigma) := \langle \hat{u} \rangle_\eta$, we end up with

$$(\partial_\sigma - \kappa_{\beta,\nu} \sigma^{\nu-1} \nabla_y^2) u(y, y', \sigma) = 0, \quad (65)$$

where $y := \sqrt{v(\sigma)}x$ actually contains a dependence on σ . Restoring the coordinates x , we finally obtain

$$[\partial_\sigma - \kappa(\sigma) \nabla_x^2] u_{\beta,\nu}(x, x', \sigma) = 0, \quad \kappa(\sigma) := \kappa_{\beta,\nu} \frac{\sigma^{\nu-1}}{v(\sigma)}. \quad (66)$$

In particular, for the fractional measure (61) $\kappa(\sigma) = \kappa_{\beta,\nu} \Gamma(\beta) \sigma^{\nu-\beta}$. Notice that the solution of the diffusion equation (66) is not the solution of (65), since the function $u[\sqrt{v(\sigma)}x, y', \sigma]$ is neither correctly normalized to 1 nor well-defined for initial points at $|x'| < \infty$. The actual fractional solution $u_{\beta,\nu}(x, x', \sigma)$ is a Gaussian PDF proportional to $u(y, y', \sigma)$:

$$u_{\beta,\nu}(x, x', \sigma) = \frac{e^{-\frac{|x-x'|^2}{4\ell^2(\sigma)}}}{[4\pi\ell^2(\sigma)]^{\frac{D}{2}}}, \quad (67)$$

where the dispersion ℓ^2 is

$$\ell^2(\sigma) := \bar{\ell}^2 + \int^\sigma d\sigma' \kappa(\sigma') = \bar{\ell}^2 + \kappa_{\beta,\nu} \int^\sigma d\sigma' \frac{\sigma'^{\nu-1}}{v(\sigma')}, \quad (68)$$

and we allowed for an additive constant $\bar{\ell}^2$. The initial condition $u_{\beta,\nu}(x, x', 0) = \delta(x - x')$ imposes that $\ell^2(0) = 0$, which fixed $\bar{\ell}^2$ depending on the measure $v(\sigma)$. In the fractional case (61), one has that $\bar{\ell} = 0$ and

$$\ell^2(\sigma) = \kappa_{\beta,\nu} \frac{\Gamma(\beta)}{1 + \nu - \beta} \sigma^{1+\nu-\beta}. \quad (69)$$

If we had set $\bar{\ell} \neq 0$, we would have obtained a Gaussian initial spread $u_{\beta,\nu}(x, x', 0) = \exp[-|x - x'|^2/(4\bar{\ell}^2)]/(4\pi\bar{\ell}^2)^{D/2}$ of width $\bar{\ell}$, a situation mimicking the idea of a particle on a fuzzy manifold with a minimal length [46, 47, 48, 49, 50, 51]. We will come back to this point in section 5, where a non-vanishing $\bar{\ell}$ is necessary in certain multi-scale scenarios.

Equation (66) is not yet in its final form, since we have not discussed the normalization of the solution in a multi-scale spacetime. In spacetimes with non-trivial measure, the normalization of the PDF is done with respect to the total measure weight $v(x)$. The only way to get the result and maintain both the Langevin-equation interpretation and the delta initial condition is to define the fractional PDF as

$$P_{\beta,\nu}(x, x', \sigma) = \frac{u_{\beta,\nu}(x, x', \sigma)}{v(x)}, \quad (70)$$

which is the solution of

$$[\partial_\sigma - \kappa(\sigma)\check{\mathcal{K}}_v]P_{\beta,\nu}(x, x', \sigma) = 0, \quad P_{\beta,\nu}(x, x', 0) = \delta_v(x, x'). \quad (71)$$

Since $u_{\beta,\nu}$ obeys the self-similarity relation

$$u_{\beta,\nu}[\lambda^{(1+\nu-\beta)/2}x, \lambda^{(1+\nu-\beta)/2}x', \lambda\sigma] = \lambda^{-D(1+\nu-\beta)/2}u_{\beta,\nu}(x, x', \sigma), \quad (72)$$

$P_{\beta,\nu}$ is self-similar with scaling law

$$P_{\beta,\nu}[\lambda^{(1+\nu-\beta)/2}x, \lambda^{(1+\nu-\beta)/2}x', \lambda\sigma] = \lambda^{-D\alpha(1+\nu-\beta)/2}P_{\beta,\nu}(x, x', \sigma). \quad (73)$$

Quantum mechanics motivates the appearance of the operator $\check{\mathcal{K}}_v$ in the diffusion equation (71) independently (appendix A). It also determines, independently from the Langevin-equation approach, the natural diffusion time (68) (appendix B).

4.1.1. Factorizable-spacetime Brownian motions Set first $\nu = 1$. From equations (65) and (62), it follows that X is the stochastic process

$$\boxed{X_{\text{FSBM}-v}(\sigma) = \frac{X_{\text{BM}}(\sigma)}{\sqrt{v(\sigma)}}.} \quad (74)$$

This is a plain *Brownian Motion*, but in a fractional or, in general, *Factorizable multi-scale Spacetime* (FSBM). We label FSBM in this model of multi-scale geometry with weighted Laplacian as FSBM- v . Some trajectories for the weight (61) are shown in figure 1. The raggedness of the curve and the drift from the average decrease when β increases.

Together with equation (5), equations (61) and (74) imply the self-similarity property

$$X_{\text{FSBM}-v}(\lambda\sigma) = \lambda^{\frac{2-\beta}{2}}X_{\text{FSBM}-v}(\sigma). \quad (75)$$

The mean squared displacement of FSBM is anomalous,

$$\langle X_{\text{FSBM}-v}^2(\sigma) \rangle = \int d^Dx v(x) P_{\beta,\nu}(x, x', \sigma) x^2 \propto \ell^2(\sigma), \quad (76)$$

and, for the fractional case $\ell^2(\sigma) \propto \sigma^{2-\beta}$, one has sub-diffusion when $0 < \beta < 1$. Moreover, increments are uncorrelated but not stationary, since their distribution does not depend on the time interval only:

$$\langle [X_{\text{FSBM}-v}(\sigma) - X_{\text{FSBM}-v}(\sigma')]^2 \rangle \neq \langle X_{\text{FSBM}-v}^2(\sigma - \sigma') \rangle. \quad (77)$$




Figure 1. Thick (black) curve: example of trajectory of the process (74) in a fractional spacetime with weighted Laplacian, diffusion measure weight (61), $\nu = 1$, and $\beta = 1/2$ (top panel), $\beta = 1$ (middle panel, ordinary Brownian motion), and $\beta = 3/2$ (bottom panel). Thin (blue) curve: example of trajectory of ordinary Brownian motion in ordinary spacetime, plotted for reference (notice the different scaling of the vertical axes).

If $\nu \neq 1$, set $\xi = \sigma^{(\nu-1)/2}\eta$, where η is a Gaussian white noise (2). Then, the Langevin equation (from (63)) $\partial_\sigma Y = \xi$ becomes

$$\partial_\sigma X_{\text{SBM}}(\sigma) = \sigma^{\frac{\nu-1}{2}}\eta(\sigma), \quad (78)$$

where we called $Y(\sigma) = X_{\text{SBM}}(\sigma)$ the solution. This is nothing but a *scaled Brownian motion* (SBM) [35, 36, 24], defined as

$$X_{\text{SBM}}(\sigma) := X_{\text{BM}}(\sigma^\nu). \quad (79)$$

Its self-similarity property is $X_{\text{SBM}}(\lambda\sigma) = \lambda^{\nu/2}X_{\text{SBM}}(\sigma)$. Just like FSBM, SBM is Markovian, since the scale transformation $\sigma \rightarrow \sigma^\nu$ preserves time ordering for $\nu > 0$ [35]. Again, increments are non-stationary, $\langle [X_{\text{SBM}}(\sigma) - X_{\text{SBM}}(\sigma')]^2 \rangle = \langle [X_{\text{BM}}(\sigma^\nu) - X_{\text{BM}}(\sigma'^\nu)]^2 \rangle = \langle X_{\text{BM}}^2(\sigma^\nu - \sigma'^\nu) \rangle \neq \langle X_{\text{SBM}}^2(\sigma - \sigma') \rangle$. Figure 2 shows the effect of the time rescaling: as ν decreases, the trajectory becomes smoother.

Equations (74) and (75) are replaced by a fractional-spacetime scaled Brownian motion (FSSBM in short):

$$X(\sigma) = X_{\text{FSSBM}}(\sigma) := \frac{X_{\text{SBM}}(\sigma)}{\sqrt{v(\sigma)}}, \quad (80)$$

with scaling law

$$X_{\text{FSSBM}}(\lambda\sigma) = \lambda^{\frac{1+\nu-\beta}{2}} X_{\text{FSSBM}-v}(\sigma). \quad (81)$$

Its variance is proportional to ℓ^2 , which for the fractional case (61) is

$$\langle X_{\text{FSSBM}}^2(\sigma) \rangle \propto \sigma^{1+\nu-\beta}. \quad (82)$$

We notice that, in an ordinary spacetime, equation (66) with time-dependent diffusion coefficient can describe two different stochastic processes. One is SBM. Another is fractional Brownian motion (FBM) [52, 53] (see also [30, 31, 33, 34, 35, 36, 37, 24]). In this case, the GLE features a Weyl fractional integral of order $\gamma = (2 + \nu - \beta)/2$ and reads $\partial_\sigma X_{\text{FBM}}(\sigma) = \partial_\sigma [-_\infty I^\gamma \eta](\sigma)$. Scaled and fractional Brownian motion are *twins* [24], i.e., they share exactly the same diffusion equation (66) and the mean squared displacement. However, they are physically quite different, since FBM neither is Markovian nor has stationary increments. From the point of view of fractional spaces, there is no twin problem and no ambiguity. In fact, the natural Langevin equation is

$$\partial_\sigma [\sqrt{v(\sigma)} X] = \xi, \quad (83)$$

which is neither the one of SBM nor of FBM. Furthermore, we are dealing with a Markovian process, as the derivation of the diffusion equation has shown. The Markovian stochastic process associated with equations (83) and (66) is unambiguously identified with FSSBM, or with FSBM- v if $\nu = 1$.




Figure 2. Thick (black) curve: example of trajectory of scaled Brownian motion (79) in ordinary spacetime with $\nu = 1/2$ (top panel), $\nu = 1$ (middle panel, ordinary Brownian motion), and $\nu = 3/2$ (bottom panel). The same trajectories represent the process (89) in a fractional spacetime with ordinary Laplacian with $\nu = 1$ and, respectively, $\beta = 3/2, 1, 1/2$. Thin (blue) curve: example of trajectory of ordinary Brownian motion in ordinary spacetime, plotted for reference (notice the different scaling of the vertical axes). The code implements the algorithm presented in equation (A2) of [35].

4.1.2. *Spectral and walk dimension* For a generic measure $v(\sigma)$, the spectral dimension is

$$\begin{aligned} d_S(\sigma) &\stackrel{(24)}{=} -2 \frac{d \ln Z(\ell)}{d \ln \ell^2} \frac{d \ln \ell^2(\sigma)}{d \ln \sigma} \\ &\stackrel{(67)}{=} D \frac{d \ln \ell^2(\sigma)}{d \ln \sigma} \end{aligned} \quad (84)$$

$$\stackrel{(68)}{=} D \frac{\sigma \kappa(\sigma)}{\int^\sigma d\sigma' \kappa(\sigma')} . \quad (85)$$

In the case (68), the heat kernel is $Z(\sigma) \propto \sigma^{-D(1+\nu-\beta)/2}$ (where the proportionality constant is the total integer volume) and

$$\boxed{d_S = D(1 + \nu - \beta) .} \quad (86)$$

The spectral dimension depends both on the choice of weight for σ and, via the parameter ν , on the statistics of the underlying stochastic process. We distinguish various cases:

- (i) A natural assumption (also supported by the findings in appendix B) is that this process is the counterpart of Brownian motion, so $\nu = 1$ and $d_S = D(2 - \beta)$.
 - (a) If one further assumes that $\beta = 1$, one recovers the case of the old diffusion equation with non-regularized heat kernel Z , section 3.2, where $d_S = D$. Diffusion is non-anomalous, $\langle X_{\text{FSSBM}}^2(\sigma) \rangle \propto \sigma$.
 - (b) Assuming that σ inherits the measure of time coordinate, $\beta = \alpha_0$, one has $d_S = D(2 - \alpha_0) \geq D$ and super-diffusion.
 - (c) Assuming instead that the charge β is equal to the average fractional charge (34), $\beta = \alpha$, $d_S = D(2 - \alpha) \geq D$ and again one has super-diffusion.
- (ii) The evolution parameter σ does not have to inherit the measure of time coordinate, and there is no compelling reason why β should coincide with the fractional charges α_0 or α , as in cases (b) and (c). One could set $\beta = 1$ from the start, as in the old diffusion equation, but reintroduce α dependence ‘from the backdoor’, via a scaled Brownian motion $X = Y$ with $\nu = \alpha$. Then $d_S = D\alpha$ as in the old diffusion equation case with regularized $\mathcal{P}$.

None of these cases corresponds to a fractal in the usual sense, since from equation (82) one sees that $\langle X_{\text{FSSBM}}^2(\sigma) \rangle \propto \sigma^{d_S/D} \neq \sigma^{d_S/d_H}$. The origin of the violation of relation (27) can be understood by revisiting the argument in section 2.3 with the form of the density of states found in section 3.2. Apart from the replacement $\rho(E) \rightarrow w(E)\rho(E)$ (which, in fact, is only a redefinition of the density and one can just put $w = 1$), the main difference is in the state counting per Hausdorff volume, which is not proportional to $d\rho(x)$ but to $d^D x$. Thus, one should replace d_H in equations (28) and (31) with the topological dimension D . Thus, for this model of fractional spacetimes the relation (27) is replaced by

$$d_W = 2 \frac{D}{d_S} . \quad (87)$$

We conclude with a remark on the various choices (i)-(ii). The diffusion equation is derived, or assumed, in a classical-mechanics context. Here, the only information about an anomalous dimension is given by the measure in the time direction, while the only information about spatial dimensions is the number $D - 1$ of particles $x^i(t)$, which is simply the topological dimension of space. From the point of view of the diffusion equation, the time direction is an external, arbitrary parameter σ . There is an intrinsic element of ambiguity in the whole construction which can be removed only by a *definition* of the statistics of the random noise. Depending on which choice is regarded as ‘fundamental’, the final output will be different. The problem in this model (and the next) of fractional spaces is that none of the above choices seems to be well motivated when looking at the analogous procedure in ordinary Euclidean space. In section 4.3 we shall consider another model where this conundrum is apparently solved.

4.2. Ordinary Laplacian

Equation (71) is not self-adjoint, which is a rather common situation for general Fokker–Planck equations [29]. However, in a quantum-gravity setting, or in any case where the diffusion equation is aimed at the determination of the spectral dimension of spacetime, one may wonder whether the physical and geometric consequences of the diffusion equation change if one takes its adjoint.[§] In the case of (71), $\mathcal{K}_v^\dagger$ is simply the ordinary Laplacian and the adjoint equation is

$$[\partial_\sigma - \kappa(\sigma)\nabla_x^2]\tilde{P}_{\beta,\nu}(x, x', \sigma) = 0, \quad \tilde{P}_{\beta,\nu}(x, x', 0) = \delta_v(x, x'). \quad (88)$$

This diffusion equation actually corresponds to the scenario with standard integer Laplacian. In that case, one can show that β is the exponent governing the power law of the quantum-mechanical time (appendix B).

The associated stochastic process is the scaled Brownian motion

$$X(\sigma) = X_{\text{FSBM}}(\sigma) := X_{\text{BM}}(\sigma^{1+\nu-\beta}).$$

(89)

In fact, in order to construct the GLE associated with this spacetime, one should account for both the differential structure (ordinary derivatives) and the fractional generalization of the source ξ . The first information suggests that the left-hand side of the Langevin equation should be of the form $\partial_\sigma X$. The second information yields, in the simplest case $\nu = 1$, $\langle \xi^2 \rangle \propto \delta_v(\sigma, \sigma')$ and $\xi(\sigma) \propto \eta(\sigma)/\sqrt{v(\sigma)}$, where η is a white noise. The result for general ν is then $\partial_\sigma X = \sigma^{(\nu-\beta)/2}\eta$ for the weight (61), which is (78) with ν replaced by $1 + \nu - \beta$. The argument is somewhat heuristic, since ordinary derivatives mix with weights in this model upon integrating by parts, and it is not obvious how a complete GLE should look like. Anyway, it yields the diffusion equation (88) and the

[§] The operation of taking the adjoint of the diffusion equation is not of clear interpretation, so even if d_S differed in mutually adjoint pairs one should not draw the conclusion that there is a problem with them.

same anomalous scaling (81) of the previous model. A trajectory of the walker is shown in figure 2.

The solution of equation (88) is again a Gaussian like (67),

$$\tilde{P}_{\beta,\nu} = C(x', \sigma) \exp \left[\frac{-|x - x'|^2}{4\ell^2(\sigma)} \right], \quad (90)$$

but with a non-standard normalization C depending on the initial point x' . This is due to the initial condition in (88), different with respect to that of equation (66). We impose

$$1 = C(x', \sigma) \int_{-\infty}^{+\infty} d^D x v(x) e^{-\frac{|x-x'|^2}{4\ell^2(\sigma)}}. \quad (91)$$

Consider the case of fixed dimensionality, $v = v_\alpha$. The integral factorizes and can be done analytically for each direction ([54], formulæ 3.462.1 and 9.240). Inverting the final result yields the normalization constant

$$C(x', \sigma) = \prod_{\mu} \left\{ \frac{\Gamma(\alpha_\mu/2)}{\Gamma(\alpha_\mu)} [2\ell(\sigma)]^{\alpha_\mu} \Phi \left[\frac{1 - \alpha_\mu}{2}; \frac{1}{2}; -\frac{|x'^\mu|^2}{4\ell^2(\sigma)} \right] \right\}^{-1}, \quad (92)$$

where Φ (also called ${}_1F_1$ or M) is Kummer's confluent hypergeometric functions of the first kind:

$$\Phi(a; b; z) := \sum_{n=0}^{+\infty} \frac{(a)_n}{(b)_n} \frac{z^n}{n!}, \quad (93)$$

where $(a)_n = \Gamma(a + n)/\Gamma(a)$ is the Pochhammer symbol. To check that the initial condition in (88) is respected, we recall the asymptotic limit

$$\Phi(a; b; z) \stackrel{z \rightarrow -\infty}{\sim} \frac{\Gamma(b)}{\Gamma(b - a)} (-z)^{-a}. \quad (94)$$

Thus,

$$C(x', \sigma) \stackrel{\sigma \rightarrow 0}{\sim} \prod_{\mu} \left[\frac{\Gamma(1/2)}{\Gamma(\alpha_\mu)} \ell^{\alpha_\mu} \left| \frac{x'^\mu}{\ell} \right|^{\alpha_\mu - 1} \right]^{-1} = \frac{1}{(4\pi\ell^2)^{D/2}} \frac{1}{v_\alpha(x')}, \quad (95)$$

and one recovers the fractional delta distribution δ_v .

Next, we calculate the asymptotic limits of the heat kernel

$$Z(\sigma) = \int d^D x v_\alpha(x) \tilde{P}_{\beta,\nu}(x, x, \sigma) = \int d^D x v_\alpha(x) C(x, \sigma). \quad (96)$$

This expression is not a power law in σ , so in general it will not give a constant spectral dimension as in the adjoint case. We estimate d_S at small diffusion scales, equation (25). Using again (95), we get $Z(\sigma) \sim \mathcal{V}(4\pi\ell^2)^{-D/2}$, where $\mathcal{V} = \int d^D x$ is the ordinary volume. Thus, one recovers equation (86).

For large σ , $\Phi \rightarrow 1$, $C(x, \sigma) \sim \prod_{\mu} [\Gamma(\alpha_\mu/2)/\Gamma(\alpha_\mu) \ell^{\alpha_\mu}(\sigma)]^{-1}$ and $Z(\sigma) \propto \mathcal{V}_H \ell^{-D\alpha}(\sigma) \propto \sigma^{-D\alpha(1+\nu-\beta)/2}$, and the spectral dimension is α times smaller than the one at small scales. This mismatch can be explained as an artifact of adopting a measure corresponding to a geometry with fixed dimensionality. Characteristic scales appear

either as topological effects (as, for instance, the curvature radius of a sphere or a torus) or in dimensional flows determined by an intrinsically multi-scale measure. Here there is no foothold where to establish a scale hierarchy, so it is more natural to take equation (25) as the correct definition of d_S when characteristic scales are not expected. Therefore, the limit $\sigma \rightarrow \infty$ of the return probability has no significance here and the result for d_S can be interpreted as in agreement with the one from the adjoint diffusion equation. Further support to this conclusion will be given in section 5.2, where we shall compute d_S in the multi-scale scenarios. In the more realistic case where d_S changes with the scale, the spectral dimension of the two models is in perfect agreement.

4.3. q -Laplacian

This scenario has a standard classical mechanics (ordinary derivatives in a certain diffusion time τ). All the results of section 2 hold with the following changes:

- (i) The random variable $X(\sigma)$ is now regarded as a composite variable $Q(\tau) = Q[X(\tau)]$. In the case of fixed dimensionality, for each direction

$$Q^\mu[X(\tau)] = \frac{\text{sgn}[X^\mu(\tau)]|X^\mu(\tau)|^{\alpha_\mu}}{\Gamma(\alpha_\mu + 1)}. \quad (97)$$

This expression is invertible, since $\text{sgn}(X) = \text{sgn}(Q)$ and $X = \text{sgn}(Q)[\Gamma(\alpha + 1)|Q|]^{1/\alpha}$. The parameter τ is by itself composite, $\tau = \varrho(\sigma)$, in particular

$$\tau = \varrho_\beta(\sigma) = \frac{\sigma^\beta}{\Gamma(\beta + 1)} \quad (98)$$

in the no-scale fractional case, where σ has the dimensionality of a length and β is a constant.

- (ii) The diffusion equation is

$$\left\{ \frac{\partial}{\partial \tau(\sigma)} - \sum_\mu \kappa_{(\mu)} \frac{\partial^2}{[\partial q(x^\mu)]^2} \right\} P = 0, \quad [\tau] = -\beta, \quad [\kappa_{(\mu)}] = \beta - 2\alpha_\mu, \quad (99)$$

where, in the general anisotropic case, the diffusion coefficient is different for each direction. One can also define the parameter

$$\ell_\mu(\sigma) := \ell_\mu[\tau(\sigma)] = \sqrt{\kappa_{(\mu)}\tau(\sigma)}, \quad [\ell_\mu] = -\alpha_\mu. \quad (100)$$

In the isotropic case ($\alpha_\mu = \alpha$), there is only one diffusion coefficient κ and one parameter $\ell(\sigma) := \sqrt{\kappa\tau(\sigma)}$ with scaling $[\ell] = -\alpha$. Then, the diffusion equation (99) can be written as

$$\left\{ \frac{\partial}{\partial \ell^2(\sigma)} - \sum_\mu \frac{\partial^2}{[\partial q(x^\mu)]^2} \right\} P = 0. \quad (101)$$

This version of the diffusion equation is also valid in the anisotropic case when the dimensionful coefficients are absorbed in the definitions of q^μ , in which case $\ell^2 = \tau$.

- (iii) Thus, as a function of the actual diffusion time σ (which is the one with respect to which one must derive the heat kernel in order to get d_S) the random variable $Q(\sigma) = Q[\tau(\sigma)]$ is a scaled Brownian motion. In the particular case of equation (97), in the first orthant one can identify the stochastic process, which we dub FSBM- q , as (up to $O(1)$ constants)

$$X_{\text{FSBM-}q}(\sigma) \sim [X_{\text{BM}}(\sigma^\beta)]^{1/\alpha} = [X_{\text{SBM}}(\sigma)]^{1/\alpha}. \quad (102)$$

The raggedness of the ordinary SBM increases with increasing β (figure 2). On the other hand, with respect to an ordinary SBM the raggedness of FSBM- q is greater ($1/\alpha > 1$) and increases for smaller α ; the drift from the average is then amplified. There is an interplay of the two effects, but the effect from α (power of Q) is greater than the one from β (scaled time), even when $\beta = \alpha$. Therefore, this incarnation of FSBM is typically more ragged than an ordinary Brownian motion; see figure 3.

- (iv) Equation (20) holds with the coordinate replacement $x \rightarrow q(x)$,

$$P(x, x', \sigma) = \prod_{\mu} \frac{e^{-\frac{|q(x^\mu) - q(x'^\mu)|^2}{4\ell_\mu^2(\sigma)}}}{\sqrt{4\pi\ell_\mu^2(\sigma)}}, \quad (103)$$

where the normalization is automatically correct.

- (v) The heat kernel is of the form (53), which for the fractional case (98) reads

$$Z(\sigma) = \frac{\mathcal{V}_H}{\prod_{\mu} \sqrt{4\pi\ell_\mu^2(\sigma)}} \propto \mathcal{V}_H \sigma^{-\frac{1}{2}D\beta}, \quad (104)$$

so that the first Seeley–DeWitt coefficient does correspond to the Hausdorff volume (this fixes a problem unsolved in the other models) and the spectral dimension is

$$d_S = D\beta. \quad (105)$$

As in the other theory, the spectral dimension only depends on the topological dimension of space and on the fractional charge associated with diffusion time. The latter stems from a classical mechanical model which knows nothing about the Hausdorff dimension of spacetime, since it only sees D motions Q^μ . Again, σ is just a parameter from the point of view of spacetime, and β does not necessarily have to be equal to the time fractional charge α_0 . To fix this arbitrariness, we can encode the information of the Hausdorff dimension in β and define it as the average fractional charge, $\beta = \alpha = d_H/D$. This way, it results that $d_S = d_H$, as in equation (52).

Even for a general β , the walk dimension is indeed the one for a fractal, since from $Q \sim \tau^{1/2}$ there follows $X \sim \sigma^{\beta/(2\alpha)}$, and equation (27) holds.




Figure 3. Thick (black) curve: example of trajectory of the process (102) for the diffusion measure weight (61), with $\beta = 1 = \alpha$ (top panel, ordinary Brownian motion in ordinary spacetime) and $\beta = 1/2 = \alpha$ (bottom panel, fractional spacetime with q -Laplacian). Thin (blue) curve: example of trajectory of ordinary Brownian motion in ordinary spacetime, plotted for reference (notice the different scaling of the vertical axes).

5. Diffusion in multi-fractional spacetimes

Assume that the distribution $v(\sigma)$ takes the same form of the multi-fractional measure weight (36),

$$v(\sigma) = v_*(\sigma) = \sum_{n=1}^N g_n \sigma^{\beta_n - 1}, \quad (106)$$

and for the sake of simplicity consider the binomial case $g_1 = \ell_*^{1-\beta_*}$, $0 < \beta_1 = \beta_* < 1$, $g_2 = 1$, and $\beta_2 = 1$:

$$v_*(\sigma) = 1 + \left(\frac{\sigma}{\ell_*} \right)^{\beta_* - 1}, \quad (107)$$

where ℓ_* is a fundamental length discriminating between infrared ($\sigma \gg \ell_*$) and ultraviolet ($\sigma \ll \ell_*$). Also fix $\nu = 1$ in the statistics of the random walker and call $\kappa_{\beta,1} = \kappa_\beta$.

5.1. Weighted Laplacian

When the spacetime measure weight $v(x)$ encodes a multi-scale geometry, the theory with weighted Laplacian does not fare well, although it can be rescued by an interesting modification of the physical interpretation.

The multi-scale version of the process (74) is obtained by using the distribution (106) or (107); we do not plot the resulting trajectories here. From now on we aim to compute the spectral dimension.

Let $1 < \beta_* < 2$. From the asymptotic behaviour of (107), one expects that in the small-scale limit the first term dominates and $d_S \sim D$, while in the large-scale limit the second term takes the lead and one recovers $d_S \sim D(2 - \beta_*) \leq D$, i.e., a reduction of the spectral dimension with the scale. This is confirmed by an explicit calculation. Plugging (107) in equation (68), we obtain

$$\ell^2(\sigma) = \bar{\ell}^2 + \int^\sigma \frac{d\sigma'}{1 + (\sigma'/\ell_*)^{\beta_*-1}} = \bar{\ell}^2 + \kappa_\beta \sigma F \left[1, \frac{1}{\beta_* - 1}; \frac{\beta_*}{\beta_* - 1}; - \left(\frac{\sigma}{\ell_*} \right)^{\beta_*-1} \right], \quad (108)$$

where we used formula 3.194.5 of [54] and $F = {}_2F_1$ is the hypergeometric function

$$F(a, b; c; z) := \sum_{n=0}^{+\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!}. \quad (109)$$

For $\sigma \ll \ell_*$, $F \rightarrow 1$ and $\ell^2(\sigma) \rightarrow \bar{\ell}^2 + \kappa_\beta \sigma$, thus fixing $\bar{\ell} = 0$. For $z \rightarrow \infty$ one must analytically continue via formula 9.132.1 of [54]:

$$F(1, b; b+1; z) = \frac{b}{b-1} \frac{1}{1-z} F \left(1, 1; 2-b; \frac{1}{1-z} \right) + \Gamma(b+1)\Gamma(1-b) \frac{1}{(-z)^b}. \quad (110)$$

For $\beta_* > 1$, $b = 1/(\beta_* - 1) > 1$, so that

$$\ell^2(\sigma) \sim \begin{cases} \kappa_\beta \sigma & (\sigma \ll \ell_*) \\ \frac{\kappa_\beta \sigma}{2 - \beta_*} \left(\frac{\sigma}{\ell_*} \right)^{1-\beta_*} & (\sigma \gg \ell_*) \end{cases}, \quad \text{for } 1 < \beta_* < 2. \quad (111)$$

From equation (85), the spectral dimension reads

$$d_S(\sigma) = \frac{D \kappa_\beta \sigma}{v_*(\sigma) \ell^2(\sigma)}. \quad (112)$$

From the asymptotic behaviour (111), it immediately follows that

$$d_S \sim \begin{cases} D & (\sigma \ll \ell_*) \\ D(2 - \beta_*) \leq D & (\sigma \gg \ell_*) \end{cases}, \quad \text{for } 1 < \beta_* < 2, \quad (113)$$

as announced.

Let us now consider the case $0 < \beta_* < 1$ ($b < -1$). Again, one expects a decrease of the spectral dimension as the scale increases, from $d_S \sim D(2 - \beta_*) \geq D$ to $d_S \sim D$. Equation (108) is valid only for $\beta_* > 1$, but we can analytically continue it under the provision that $\ell^2(0) = 0$. At large scales, $\ell^2(\sigma) \sim \bar{\ell}^2 + \kappa_\beta \sigma$, while for $\sigma \gg \ell_*$ we use

equation (110), this time picking both terms. Noting that $\Gamma(b)\Gamma(1-b) = \pi/\sin(\pi b)$, from equation (108) we get

$$\ell^2(\sigma) \stackrel{\sigma \lesssim \ell_*}{\sim} \left[\bar{\ell}^2 + \kappa_\beta \ell_* \frac{\frac{\pi}{\beta_*-1}}{\sin \frac{\pi}{\beta_*-1}} \right] + \frac{\kappa_\beta \sigma}{2 - \beta_*} \left(\frac{\sigma}{\ell_*} \right)^{1-\beta_*}, \quad (114)$$

implying that

$$\bar{\ell}^2 = -\kappa_\beta \ell_* \frac{\frac{\pi}{\beta_*-1}}{\sin \frac{\pi}{\beta_*-1}}. \quad (115)$$

This formula is valid also for $\beta_* = 1 + 1/k$, $k \in \mathbb{Z}$, where $\bar{\ell}^2 \rightarrow -\kappa_\beta \ell_*$. To summarize,

$$\ell^2(\sigma) \sim \begin{cases} \frac{\kappa_\beta \sigma}{2 - \beta_*} \left(\frac{\sigma}{\ell_*} \right)^{1-\beta_*} & (\sigma \ll \ell_*) \\ \bar{\ell}^2 + \kappa_\beta \sigma & (\sigma \gg \ell_*) \end{cases}, \quad \text{for } 0 < \beta_* < 1, \quad (116)$$

and we get an interchange of the regimes of (113):

$$d_S \sim \begin{cases} D(2 - \beta_*) \geq D & (\sigma \ll \ell_*) \\ D & (\sigma \gg \ell_*) \end{cases}, \quad \text{for } 0 < \beta_* < 1, \quad (117)$$

although the decreasing behaviour remains.

The overall scenario is radically different from the ‘embedding picture’ previously associated with models with weighted Laplacians. In that case, one started with a D -dimensional ambient Minkowski spacetime M^D and had a multi-fractional spacetime $\mathcal{M}_*$ embedded in it. Then, $\mathcal{M}_*$ ‘filled’ M^D only at large scales. This is true also in the present case, but only for the Hausdorff dimension. On the contrary, there is a reduction of the spectral dimension at large scales:

- I. For $0 < \beta_* < 1$ (equation (117)), it is as if fractional spacetime gushed out of the embedding at smaller and smaller scales. Taking, for instance, $D = 4$ and $\beta = 1/4, 1/2, 3/4$, one would have a dimensional reduction $d_S \sim 7, 6, 5 \rightarrow d_S \sim 4$.
- II. For $1 < \beta_* < 2$ (equation (113)), it is as if the embedding were depleted of spacetime points at larger and larger scales. For example, for $D \geq 5$ and $\beta = 2 - 4/D$, one gets $d_S \sim 4$ in the infrared starting from an embedding with 5 or more dimensions.

Possible modifications may come from taking a more general statistics for the diffusion process ($\nu \neq 1$, but this is not natural from the point of view of quantum mechanics; see the appendices) or from changing the initial condition from a delta to a Gaussian. As discussed in section 4.1, this can be achieved simply by letting the constant $\bar{\ell}^2$ to be non-vanishing by default. The interpretation then would be the one of [46, 47, 48, 50], namely, that the test particle is not pointwise due to the intrinsic ‘fuzziness’ of a spacetime with a minimal length. To get this type of structure in the multi-fractional context, it would be quite natural to identify the characteristic scale ℓ_* in the binomial measure with the scale at which ‘fuzziness’ effects become important. In turn, this univocally sets

$$\bar{\ell} = \ell_*. \quad (118)$$

This would ‘straighten’ dimensional flow. Take, in fact, $0 < \beta_* < 1$. The only but crucial modification to the above calculation of d_S is that the constant term in equation (114) does *not* vanish. Call L^2 the content in square brackets. While the large-scale limit of the spectral dimension is unaffected, the small-scale one is

$$d_S \stackrel{\sigma \ll \ell_*}{\sim} D \frac{\kappa_\beta \ell_*}{L^2} \left(\frac{\sigma}{\ell_*} \right)^{2-\beta_*}, \quad L^2 = \bar{\ell}^2 + \kappa_\beta \ell_* \frac{\frac{\pi}{\beta_*-1}}{\sin \frac{\pi}{\beta_*-1}}, \quad (119)$$

leading to

$$d_S \sim \begin{cases} 0 & (\sigma \ll \ell_*) \\ D & (\sigma \gg \ell_*) \end{cases}. \quad (120)$$

Such a scenario somewhat merges multi-fractal^{||} and effective non-commutative geometries, but in a way different from [11]. In [11], κ -Minkowski spacetime was reinterpreted as the ultra-microscopic limit of a log-oscillating fractional measure. On the other hand, here we have changed the initial condition of diffusion (but, implicitly, also of the Green function) so that to have the scale in the real-order multi-fractional measure play the role of a fuzziness length, just as in commutative effective spacetimes coming from a non-commutative geometry where operators are evaluated on coherent states [46, 47, 48]. However, the matching is only at the level of the spectral dimension [50], since our resolution of the identity is not a Gaussian but a fractional delta. We leave a further elaboration of this scenario and a fuller assessment of its physical consequences for future study.

5.2. Ordinary Laplacian

The normalization condition (91) now features a multi-scale measure weight $v(x) = v_*(x)$. With the binomial measure (37) and in the isotropic case $\alpha_\mu = \alpha_*$, one has

$$C(x', \sigma) = \left\{ [4\pi \ell^2(\sigma)]^{\frac{D}{2}} + \ell_*^D \left[\frac{\Gamma(\alpha_*/2)}{\Gamma(\alpha_*)} \frac{2^{\alpha_*} \ell^{\alpha_*}(\sigma)}{\ell_*^{\alpha_*}} \right]^D \prod_\mu \Phi \left[\frac{1-\alpha_*}{2}; \frac{1}{2}; -\frac{(x'^\mu)^2}{4\ell^2(\sigma)} \right] \right\}^{-1}, \quad (121)$$

where $\ell^2(\sigma)$ has been computed in the previous sub-section. As before, we focus on the asymptotic limits of the heat kernel $Z(\sigma) = \int d^D x v_*(x) C(x, \sigma)$ and of the spectral dimension. For $0 < \beta_* < 2$, $\ell^2 \rightarrow 0$ when $\sigma \rightarrow 0$, and the initial condition is recovered, since from equation (94)

$$C(x', \sigma) \stackrel{\sigma \rightarrow 0}{\sim} \left[(4\pi \ell^2)^{\frac{D}{2}} v_*(x') \right]^{-1}. \quad (122)$$

Thus, $Z \sim [\ell(\sigma)]^{-D}$. Still for the whole range $0 < \beta_* < 2$, when $\sigma \rightarrow +\infty$ one has $\ell^2 \rightarrow +\infty$, $\Phi \rightarrow 1$, and again $Z \sim [\ell(\sigma)]^{-D}$. From equations (111) and (116), one eventually gets equations (113) and (117).

This result amends the incomplete discussion in section 4.2. With respect to dimensional flow, this model falls in the same class of its Hermitian-conjugate dual, i.e., the model with weighted Laplacian.

^{||} Here we can talk about fractals because $d_S \leq d_H$.

5.3. q -Laplacian

The case of q -theory with scale-dependent dimension is straightforward. One should simply replace $\ell^2(\sigma)$ in equation (101) and the distributions $q(x^\mu)$ with a generic functional of, respectively, σ and x^μ . The multi-fractional measure weight (36) determines the multi-scale form of q :

$$q(x^\mu) = \varrho_*(x^\mu) = \sum_{n=1}^N g_{\mu,n} \text{sgn}(x^\mu) |x^\mu|^{\alpha_\mu}, \quad (123)$$

where we absorbed Γ factors into the $g_{\mu,n}$'s. This also suggests the form of ℓ^2 :

$$\ell^2(\sigma) = \varrho_*(\sigma) = \sum_{n=1}^N g_n \sigma^{\beta_n}. \quad (124)$$

The simplest case is the binomial measure (107),

$$\ell^2(\sigma) = \kappa \ell_* \left[\frac{\sigma}{\ell_*} + \left(\frac{\sigma}{\ell_*} \right)^{\beta_*} \right]. \quad (125)$$

Since $0 < \beta_* < 1$ in this model, at small scales $\sigma/\ell_* \ll 1$ the second term dominates and $\ell^2 \propto \sigma^{\beta_*}$, while at large scales $\sigma/\ell_* \gg 1$ the variance of the PDF is the usual one, $\ell^2 = \kappa \sigma$.

In general, the coefficients g_n may also depend on σ , according to the type of profile one wishes to describe. Barring this possibility, from equations (101), (124), and (84), the spectral dimension reads

$$d_S(\sigma) = D \frac{\sum_n g_n \beta_n \sigma^{\beta_n}}{\sum_n g_n \sigma^{\beta_n}}. \quad (126)$$

In particular, for the case of fixed dimensionality ($N = 1$, $\beta_1 = \beta$), the spectral dimension coincides with equation (105), while for the binomial measure one has

$$d_S(\sigma) = D \frac{1 + \beta_*(\ell_*/\sigma)^{1-\beta_*}}{1 + (\ell_*/\sigma)^{1-\beta_*}}. \quad (127)$$

As anticipated, $d_S \sim D\beta_*$ at small scales $\sigma/\ell_* \ll 1$ and $d_S \sim D$ at large scales $\sigma/\ell_* \gg 1$.

Since equation (123) is not invertible, knowing the statistics of Q does not immediately lead to a statistics for X . One can, however, apply equation (102) at various asymptotic regimes with different α , and give a different, patch-wise stochastic description of spacetime for each scale range where $d_S \sim \text{const}$.

6. Discussion

The three classes of no-scale spacetime models analyzed here display constant anomalous dimensions and are associated with highly non-trivial stochastic processes, all related (via various rescalings and mappings) to Brownian motion or scaled Brownian motion. Table 1 summarizes these findings. To the best of our knowledge, these processes do

Model (symmetries of $\mathcal{L}$)	d'Alembertian	Stochastic process	d_H	d_S
Weighted Poincaré	$\frac{1}{\sqrt{v}}\square_x(\sqrt{v}\cdot)$ eq. (38)	$X_{\text{FSBM}-v}(\sigma)$ eq. (74)	$D\alpha$ eq. (35)	$D(2-\beta)$ eq. (86)
Ordinary Poincaré	$\square_x, \square_x^\dagger = \frac{1}{v}\square_x(v\cdot)$ eqs. (43) and (44)	$X_{\text{FSBM}}(\sigma)$ eq. (89)	$D\alpha$	$D(2-\beta)$
q -Poincaré	$\square_{q(x)}$ eq. (46)	$X_{\text{FSBM}-q}(\sigma)$ eq. (102)	$D\alpha$	$D\beta$ eq. (105)

Table 1. Models of fractional spacetimes with fixed dimensionality and $\nu = 1$, characterized by the symmetries of the Lagrangian density, a stochastic process $X(\sigma)$, Hausdorff dimension d_H and spectral dimension d_S .

not have counterparts in the literature of probability theory and we had to name them according to the spacetime they describe.

We also got the analytic expression of the spectral dimension of the multi-scale extensions of these theories, found that the models with weighted and ordinary Laplacian (which are Hermitian conjugate) produce the same dimensional flow, and given an alternative interpretation of the theory with weighted Laplacian as a fuzzy spacetime. In this case, dimensional flow is modified accordingly and $d_S \rightarrow 0$ in the UV.

Only the model with q -Laplacian is fractal in the usual sense. Namely, the first Seeley–DeWitt coefficient in the heat kernel expansion corresponds to the Hausdorff volume and the relation between Hausdorff, spectral, and walk dimension is (27). The other models violate these conditions, mainly because the effective density of states scales as the integer embedding volume. Such properties are likely to signal a difference in the renormalization of field theories living on these spacetimes.

In all the models of fractional and multi-scale spacetimes we have considered in this paper, the spectral dimension was derived starting from statistical mechanics, via a generalized Langevin equation. There is, in general, an intrinsic ambiguity in the diffusion equation approach, inasmuch as it does not fix the scaling dimension of effective diffusion time. This, in fact, is part of the definition of the theory, and one should not expect to get a certain value of d_S without imposing a certain number of defining conditions. Part of this ambiguity might be related to other aspects of the theory which we have not considered here, such as the momentum-space structure. Taking as a guiding principle the conjecture [45] according to which the spectral dimension d_S is the dimension of momentum space, and fixing the latter by assuming that momentum transform is an automorphism (hence, $w(k) = v(k)$), one would fix the value of β a posteriori. Similar considerations are not rigorous at the present stage and further study will be needed.

The above approach is classical. Quantum mechanics can provide valuable information about the diffusion equation, which, as a matter of fact, can be derived as the classical limit of a quantum diffusion process. In the appendices we discuss some

aspects of the relation between classical and quantum diffusion.

Acknowledgments

The work of G C is under a Ramón y Cajal tenure-track contract.

A. Quantum probability density as a bilinear

In finding the diffusion equation (71) we did not give a robust motivation to the form of the Laplacian $\hat{\mathcal{K}}_v$, i.e., of the functional form (70). We can do so by a simple calculation of the quantum probability density function in multi-scale spacetimes with weighted Poincaré symmetries, closely following the same procedure as in ordinary spacetime [55]. Since we are in a quantum-mechanical context, we denote time with t . We also write D -dimensional spatial vectors in normal font.

Consider the quantum-mechanical free particle with unit mass and Hamiltonian $\hat{H}_{\text{free}} = -\mathcal{K}_v^E$, where $\mathcal{K}_v^E$ is the Euclidean version of (38) in D spatial dimensions. The energy $E_k = k^2$ are given by the eigenvalue equation $\hat{H}_{\text{free}}\mathfrak{e}(k, x) = E_k\mathfrak{e}(k, x)$, where we used the ‘plane waves’ (39). The associated Green equation with delta source both in space and time ($\hbar = 1$) is

$$(i\mathcal{D}_t - \hat{H}_{\text{free}})G_{\text{free}}(x, t; x', t') = \delta_{v_0}(t, t') \delta_v(x, x'). \quad (128)$$

The solutions are the advanced and retarded propagators

$$G_{\text{free}}^{\text{R,A}}(x, t; x', t') = \mp i \frac{\theta[\pm(t - t')]}{\sqrt{v_0(t)v_0(t')}} \langle x | e^{-i\hat{H}_{\text{free}}(t-t')} | x' \rangle, \quad (129)$$

where θ is the Heaviside step function. Transforming into energy-momentum space, and using

$$\mp i\theta(\pm z) = \int_{-\infty}^{+\infty} d\omega \frac{e^{i\omega z}}{\omega \pm i\epsilon}, \quad (130)$$

it is easy to show that,

$$G_{\text{free}}^{\text{R,A}}(x, t; x', t') = \int dE w(E) \mathfrak{e}^*(E, t) \mathfrak{e}(E, t') \tilde{G}_{\text{free}}^{\text{R,A}}(x, x'; E), \quad (131)$$

where we assumed a generic measure weight $w(E)$, we extended the ‘plane waves’ (39) to the energy-time pair, $\mathfrak{e}(E, t) = e^{iEt}/\sqrt{v_0(t)w(E)}$, and

$$\tilde{G}_{\text{free}}^{\text{R,A}}(x, x'; E) = \int d^D k w(k) \frac{\mathfrak{e}^*(k, x) \mathfrak{e}(k, x')}{E - E_k \pm i\epsilon}. \quad (132)$$

Making measure factors explicit and cancelling them, we obtain that the fractional propagators $G_{\text{free}}^{\text{R,A}}$ are related to the ones in ordinary space $\bar{G}_{\text{free}}^{\text{R,A}}$ by

$$G_{\text{free}}^{\text{R,A}}(x, t; x', t') = \frac{\bar{G}_{\text{free}}^{\text{R,A}}(x - x', t - t')}{\sqrt{v(x)v_0(t)v(x')v_0(t')}}. \quad (133)$$

Comparing with the propagator computed in [14], there is an extra prefactor $1/v_0(t')$ (there, the overall time-dependent prefactor is $\sqrt{v_0(t')/v_0(t)}$), because the propagator

of [14] was solution of the Green equation only with spatial delta source. Related to this, notice that, in the limit $t \rightarrow t'$, $G_{\text{free}}^{\text{R,A}}(x, t; x', t) = \delta_v(x, x')/v_0(t)$. This suggests, as a feedback, to define the Green equation (128) with an ordinary time direction (ordinary delta and time derivative); its solutions are then $\sqrt{v_0(t')v_0(t)}G_{\text{free}}^{\text{R,A}}$.

Let us now turn to a generic quantum system without specifying the Hamiltonian $\hat{H}$ and the dispersion relation between the energy E_k and momentum. We only assume that $\hat{H}$ is characterized by energy eigenvalues E_k and eigenstates $|\psi_k\rangle$: $\hat{H}|\psi_k\rangle = E_k|\psi_k\rangle$. Here the momentum k can be either discrete [55] or continuous, but for simplicity we stick with the continuum case. Let $|\Psi_{x'}\rangle$ be a normalizable wave packet of average energy $\mathcal{E}$ centered at the spatial point x' and with dispersion s :

$$|\Psi_{x'}\rangle := A \int d^D k w(k) \langle \psi_k | x' \rangle e^{-\frac{(E_k - \mathcal{E})^2}{4s}} |\psi_k\rangle, \quad (134)$$

where $A = A(x', s)$ is a normalization constant and we assumed a Gaussian form for the packet. Projecting on the position-space basis and defining $\psi_k(x) := \langle x | \psi_k \rangle$, one obtains the wave-function

$$\Psi_{x'}(x) := \langle x | \Psi_{x'} \rangle = A \int d^D k w(k) \psi_k^*(x') \psi_k(x) e^{-\frac{(E_k - \mathcal{E})^2}{4s}}. \quad (135)$$

Scalar products of wave-functions include a non-trivial weight factor, so that the orthonormality of the $\psi_k(x)$ is

$$\langle \psi_{k'} | \psi_k \rangle := \int d^D x v(x) \psi_k(x) \psi_{k'}^*(x) = \delta_w(k, k'). \quad (136)$$

This relation is the starting point for finding the normalization constant A by imposing $\langle \Psi_{x'} | \Psi_{x'} \rangle = 1$. Defining the density of states (54), one has

$$1 = A^2 \int dE w(E) \rho_{x'}(E) e^{-\frac{(E - \mathcal{E})^2}{2s}}, \quad (137)$$

where we made explicit the dependence of ρ on x' via a subscript. The result for A depends on the way the density of states is approximated. For instance, since $\rho_{x'}(E) \propto |\psi_k(x')|^2 \propto [v(x')]^{-1}$, it is natural to replace $\rho_{x'}(E)$ with $\rho_0/v(x')$, where ρ_0 is the constant average over disorder per integer volume [55]. Thus, we obtain that

$$A^2 = \frac{v(x')}{\rho_0} \mathcal{C}(s, \mathcal{E}), \quad (138)$$

where the function $\mathcal{C}$ depends on the form of $w(E)$ and, in fractional spaces, also on the average energy $\mathcal{E}$. The details of $\mathcal{C}$ are not important here, since we wish to extract only the measure-weight dependence of the quantum PDF.

The latter is defined as the square of the matrix element of the evolution operator

$$\mathcal{G}^{\text{R}}(x, t; x', t') := -i \frac{\theta(t - t')}{\sqrt{v_0(t)v_0(t')}} \langle x | e^{-i\hat{H}(t-t')} | \Psi_{x'} \rangle \quad (139)$$

$$= \int d^D y v(y) G^{\text{R}}(x, t; y, t') \Psi_{x'}(y), \quad (140)$$

averaged over disorder. The Hamiltonian, in fact, may be quite complicated by the interaction of the particle with the underlying medium. This interaction is of stochastic

type and may be included as an effective noise potential. We denote this stochastic average as $\langle\langle\cdot\rangle\rangle$. Thus, the probability density function of *quantum* diffusion is

$$\begin{aligned}
P_q(x, t; x', t') &:= \langle\langle |\mathcal{G}^R(x, t; x', t')|^2 \rangle\rangle \\
&= A^2 \int dE w(E) e^*(E, t) e(E, t') \int dE' w(E') e^*(E', t') e(E', t) \\
&\quad \times e^{-\frac{(E-\varepsilon)^2 + (E'-\varepsilon)^2}{4s}} \langle\langle \tilde{G}^R(x, x'; E) \tilde{G}^A(x', x; E') \rangle\rangle \\
&= \frac{A^2}{v_0(t)v_0(t')} \int dE \int dE' e^{-i(E-E')(t-t')} e^{-\frac{(E-\varepsilon)^2 + (E'-\varepsilon)^2}{4s}} \\
&\quad \times \langle\langle \tilde{G}^R(x, x'; E) \tilde{G}^A(x', x; E') \rangle\rangle \\
&\stackrel{\omega = \frac{E-E'}{2}}{=} \frac{A^2}{v_0(t)v_0(t')} \int dE \int d\omega e^{-i\omega(t-t')} e^{-\frac{(E-\varepsilon)^2 + (E-\omega-\varepsilon)^2}{4s}} \\
&\quad \times \langle\langle \tilde{G}^R(x, x'; E) \tilde{G}^A(x', x; E - \omega) \rangle\rangle, \tag{141}
\end{aligned}$$

where we used the analogue of equation (131) for the interacting case. At this point, we make two assumptions [55]: first, that $\omega \ll s$ (spread much larger than the frequency) and, second, that

$$P_q(x, x'; \omega) := \frac{1}{\rho_0} \langle\langle \tilde{G}^R(x, x'; E) \tilde{G}^A(x', x; E - \omega) \rangle\rangle \tag{142}$$

depends on the energy E only weakly. We can thus take the stochastic average outside of the integral in E . Then, equations (138) and (141) yield

$$P_q(x, t; x', t') \approx \frac{v(x')}{v_0(t')v_0(t)} \int d\omega e^{-i\omega(t-t')} P_q(x, x'; \omega). \tag{143}$$

The extra prefactor $1/[v_0(t')v_0(t)]$ would disappear if one assumed a Green equation with ordinary time delta and time derivative instead of (128).

From equations (133), (142) and (143), it follows that the coordinate dependence of the quantum PDF in fractional spaces is the usual one times a prefactor (spatial vector notation reinstated) $v(\mathbf{x}')[v(\mathbf{x}')v(\mathbf{x})v_0(t')v_0(t)]^{-1} = [v(\mathbf{x})v_0(t)]^{-1}[v_0(t')]^{-1} \propto [v(x)]^{-1}$. The *classical* PDF stems from various approximations of P_q [55], which however do not alter this prefactor. This is the origin of the functional form of equation (70).

B. Quantum-mechanical time

In the previous section, we derived the PDF of a quantum diffusive process and found that, as in the standard case [55], it is of the form $P_q \sim GG$, where G is the propagator of the particle. The classical PDF P can then be derived from P_q . (It is too heuristic to regard the diffusion equation as the Wick-rotated version of the Schrödinger equation and the classical PDF P as the Euclideanized version of the quantum propagator G .) We thus obtained the coordinate-dependent normalization (70). This is not the only information one can extract from quantum mechanics. In fact, the time-space dependence of the propagator G , calculated from the path integral, fixes the natural quantum-mechanical time T , i.e., the time with respect to which the path integral yields

the transition probability. Since the scaling ratio of this time with respect to spatial coordinates filters down to the classical level, this provides an independent identification of diffusion time and a check of the diffusion equations (71) and (88) in the fixed-dimensionality case, where the relation between dispersion and diffusion time is given by equation (68). To this purpose, the calculation of the free-particle propagator G_{free} suffices. In what follows, the symbol q always denotes canonical coordinates (one spatial dimension for simplicity), while for geometric coordinates we reserve the symbol $\varrho(x)$.

B.1. Weighted Lagrangian

The quantum mechanics corresponding to the fractional scenarios with weighted Laplacians features derivatives $\mathcal{D}$ in the Lagrangian. The free-particle action is

$$S = \int_{t'}^t dt'' v_0(t'') L, \quad (144)$$

where $L = \frac{1}{2}m(\mathcal{D}_t q)^2$. The Green function as a path integral

$$G \sim \sum_{\text{paths}} e^{iS} \quad (145)$$

was computed in [14] for this system and it reads

$$G_{\text{free}}(q, t; q', t') \sqrt{\frac{v_0(t')}{v_0(t)v(q)v(q')}} \sqrt{\frac{im}{2\pi(t-t')}} \exp \left\{ \frac{im[\sqrt{v_0(t)}q - \sqrt{v_0(t')}q']^2}{2(t-t')} \right\}. \quad (146)$$

The relative scaling between space and time coordinates thus roughly identifies the quantum-mechanical time

$$T \sim \frac{t}{v_0(t)}, \quad (147)$$

which, for a power-law measure, yields

$$T \propto t^{2-\beta}, \quad (148)$$

in agreement with (69) in the case $\nu = 1$. Equation (147) is not a rigorous definition stemming from equation (146), where the time-space scaling ratio is quite entangled. This is the reason why we do not get a precise matching with equation (68), which instead we obtain in the next example.

B.2. Ordinary Lagrangian

The standard-Laplacian scenario corresponds to a Lagrangian $L = \frac{1}{2}m\dot{q}^2$. For this case we report the full calculation, which is similar to the one in [14]. Partitioning the time interval $t - t'$ into N infinitesimal parts, $t = t_N > t_{N-1} > \dots > t_1 > t_0 = t'$, the action is

$$S = \frac{m}{2} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} dt v_0(t) [\dot{q}(t)]^2 \approx \frac{m}{2} \sum_{n=0}^{N-1} v_0(t_{n+1}) \frac{[q(t_{n+1}) - q(t_n)]^2}{t_{n+1} - t_n}, \quad (149)$$

where in the second step one could replace $v_0(t_{n+1})$ with $v_0(t_n)$ or the average $[v_0(t_{n+1}) - v_0(t_n)]/2$ without changing the final result.

To sum over all trajectories and get (145), it is sufficient to integrate over all possible $q_n = q(t_n)$ (ordinary integration, without weights):

$$G_N := K_N \int dq_1 \dots dq_{N-1} \exp \left[-\frac{m}{2i\epsilon} \sum_{n=0}^{N-1} v_0(t_{n+1})(q_{n+1} - q_n)^2 \right], \quad (150)$$

where K_N is a constant and we chose a partition with identical segments $\epsilon = t_{n+1} - t_n$, $N\epsilon = t - t'$. It is easy to show by iteration that

$$G_N(q_N, t_N; q_0, t_0) = K_N \prod_{n=1}^N \sqrt{\frac{2\pi i \epsilon}{m v_0(t_n)}} \sqrt{\frac{m}{2\pi i} \frac{1}{T_N}} \exp \left[-\frac{m}{2i} \frac{(q_N - q_0)^2}{T_N} \right], \quad (151)$$

where

$$T_N := \sum_{n=1}^N \frac{\epsilon}{v_0(t_n)}. \quad (152)$$

The constant K_N can be chosen so that $G_N(q_N, q_0) = \int dq_M G_N(q_N, q_M) G_M(q_M, q_0)$, which implies $K_N = K_M K_{N-M}$. Noticing that this relation is satisfied by

$$K_N = \prod_{n=1}^N \sqrt{\frac{m v_0(t_n)}{2\pi i \epsilon}}, \quad (153)$$

we thus have

$$G_N(q_N, t_N; q_0, t_0) = \sqrt{\frac{m}{2\pi i} \frac{1}{T_N}} \exp \left[-\frac{m}{2i} \frac{(q_N - q_0)^2}{T_N} \right]. \quad (154)$$

In the double limit $N \rightarrow \infty$, $\epsilon \rightarrow 0$, with $N\epsilon = t - t'$ fixed, we finally get ($q_N = q$, $q_0 = q'$)

$$\begin{aligned} G_{\text{free}}(q, t; q', t') &:= \lim_{N \rightarrow \infty} G_N(q, t; q', t') \\ &= \sqrt{\frac{m}{2\pi i} \frac{1}{T}} \exp \left[-\frac{m}{2i} \frac{(q - q')^2}{T} \right], \end{aligned} \quad (155)$$

where

$$T = \lim_{N \rightarrow \infty} T_N = \int_{t'}^t \frac{dt''}{v_0(t'')}, \quad (156)$$

which coincides with equation (68) with $\nu = 1$. For a power-law weight $v_0(t) \propto t^{\beta-1}$, equations (156) and (147) coincide with (148) up to a positive constant, but otherwise they are different.

B.3. q -Lagrangian

This case is obvious, and from the ordinary Green function it follows that $T = \varrho(t)$. In the case of fixed dimensionality, $T \propto t^\beta$.

- [2] Rammal R and Toulouse G 1983 *J. Phys. Lett. (Paris)* **44** 13
- [3] Watanabe H 1985 *J. Phys. A* **18** 2807
- [4] Havlin S and Ben-Avraham D 1987 *Adv. Phys.* **36** 695
- [5] Kigami J and Lapidus M L 1993 *Commun. Math. Phys.* **158** 93
- [6] ben-Avraham D and Havlin S 2000 *Diffusion and Reactions in Fractals and Disordered Systems* (Cambridge: Cambridge University Press)
- [7] Calcagni G 2011 *Phys. Rev. D* **84** 061501(R) ([arXiv:1106.0295](#))
- [8] Calcagni G 2012 *Adv. Theor. Math. Phys.* **16** 549 ([arXiv:1106.5787](#))
- [9] Calcagni G 2012 *J. High Energy Phys.* JHEP01(2012)065 ([arXiv:1107.5041](#))
- [10] Calcagni G and Nardelli G 2012 *Adv. Theor. Math. Phys.* **16** in press ([arXiv:1202.5383](#))
- [11] Arzano M, Calcagni G, Oriti D and Scalisi M 2011 *Phys. Rev. D* **84** 125002 ([arXiv:1107.5308](#))
- [12] Calcagni G 2012 *Phys. Rev. D* **86** 044021 ([arXiv:1204.2550](#))
- [13] Calcagni G 2013 *Phys. Rev. E* **87** 012123 ([arXiv:1205.5046](#))
- [14] Calcagni G, Nardelli G and Scalisi M 2012 *J. Math. Phys.* **53** 102110 ([arXiv:1207.4473](#))
- [15] Calcagni G 2012 ([arXiv:1209.4376](#))
- [16] Calcagni G and Nardelli G 2013 *Phys. Rev. D* **87** 085008 ([arXiv:1210.2754](#))
- [17] Calcagni G 2012 *AIP Conf. Proc.* **1483** 31 ([arXiv:1209.1110](#))
- [18] Calcagni G 2010 *Phys. Rev. Lett.* **104** 251301 ([arXiv:0912.3142](#))
- [19] Calcagni G 2010 *J. High Energy Phys.* JHEP03(2010)120 ([arXiv:1001.0571](#))
- [20] Calcagni G 2011 *Phys. Lett. B* **697** 251 ([arXiv:1012.1244](#))
- [21] S Carlip 2009 *AIP Conf. Proc.* **1196** 72 ([arXiv:0909.3329](#))
- [22] Sotiriou T P, Visser M and Weinfurtner S 2011 *Phys. Rev. D* **84** 104018 ([arXiv:1105.6098](#))
- [23] Calcagni G, Eichhorn A and Saueressig F 2013 in progress
- [24] Sokolov I M 2012 *Soft Matter* **8** 9043
- [25] Fick A 1855 *Pogg. Ann. Phys. Chem.* **170** 59
- [26] Einstein A 1905 *Annalen Phys.* **322** 549
- [27] von Smoluchowski M 1906 *Annalen Phys.* **326** 756
- [28] Kubo R, Toda M and Hashitsume N 1985 *Statistical Physics II – Nonequilibrium Statistical Mechanics* (Berlin: Springer-Verlag)
- [29] Zwanzig R 2001 *Nonequilibrium Statistical Mechanics* (Oxford: Oxford University Press)
- [30] Wang K G and Lung C W 1990 *Phys. Lett. A* **151** 119
- [31] Wang K G 1992 *Phys. Rev. A* **45** 833
- [32] Masoliver J and Wang K G 1995 *Phys. Rev. E* **51** 2987
- [33] Wang K G and Tokuyama M 1999 *Physica A* **265** 341
- [34] Lutz E 2001 *Phys. Rev. E* **64** 051106
- [35] Lim S C and Muniandy S V 2002 *Phys. Rev. E* **66** 021114
- [36] Metzler E and Klafter J 2004 *J. Phys. A* **37** R161
- [37] Jeon J-H and Metzler R 2010 *Phys. Rev. E* **81** 021103 ([arXiv:1001.0681](#))
- [38] Metzler R and Klafter J 2000 *Phys. Rept.* **339** 1
- [39] Zaslavsky G M 2002 *Phys. Rept.* **371** 461
- [40] Avramidi I G 2000 *Heat Kernel and Quantum Gravity* (Berlin: Springer-Verlag)
- [41] Kirsten K 2001 *Spectral Functions in Mathematics and Physics* (Boca Raton: Chapman & Hall/CRC)
- [42] Vassilevich D V 2003 *Phys. Rept.* **388** 279 ([arXiv:hep-th/0306138](#))
- [43] Nigmatullin R R and Le Méhauté A 2005 *J. Non-Cryst. Solids* **351** 2888
- [44] Akkermans E 2012 ([arXiv:1210.6763](#))
- [45] Akkermans E, Dunne G V and Teplyaev A 2010 *Phys. Rev. Lett.* **105** 230407 ([arXiv:1010.1148](#))
- [46] Smailagic A and Spallucci E 2003 *J. Phys. A* **36** L467 ([arXiv:hep-th/0307217](#))
- [47] Smailagic A and Spallucci E 2003 *J. Phys. A* **36** L517 ([arXiv:hep-th/0308193](#))
- [48] Smailagic A and Spallucci E (*A* **37** **7169** 2004 [arXiv:hep-th/0406174](#))
- [49] Spallucci E, Smailagic A and Nicolini P 2006 *Phys. Rev. D* **73** 084004 ([arXiv:hep-th/0604094](#))

- [50] Modesto L and Nicolini P 2010 *Phys. Rev. D* **81** 104040 ([arXiv:0912.0220](#))
- [51] Nicolini P and Niedner B 2011 *Phys. Rev. D* **83** 024017 ([arXiv:1009.3267](#))
- [52] Barnes J A and Allan D W 1966 *Proc. IEEE* **54** 176
- [53] Mandelbrot B B and Van Ness J W 1968 *SIAM Rev.* **10** 422
- [54] Gradshteyn I S and Ryzhik I M 2000 *Table of Integrals, Series, and Products* (London: Academic Press, London)
- [55] Akkermans E and Montambaux G 2011 *Mesoscopic Physics of Electrons and Photons* (Cambridge: Cambridge University Press)